

Informe Técnico realizado por Investigadores de la Red de Estudios Ambientales Bonaerense (CONICET) sobre la Modificación de la Resolución Presidencial ACUMAR N° 46/2017 y su Reglamentación a solicitud del Ministerio Público Fiscal.

10 de Julio de 2019

Coordinación científica: Dr. Jorge Sambeth sambethje@gmail.com

Colaboradores: Dr. Jorge Herkovits(herkovits.j@gmail.com), Dr. Andres Porta (aaporta@yahoo.com.ar), Dr. Pablo Demetrio (pablo.demetrio@gmail.com), Dr. Andres Arias (aharias@criba.edu.ar), Dr. Jorge Marcovecchio (jmjorgemar@gmail.com), Dra. Sandra Botté (sbotte@iado-conicet.gob.ar), Dr. Marcelo Pereyra (mpereyra@criba.edu.ar), Lic. Lautaro Girones (gironeslautaro@gmail.com), Dra. Noelia La Colla (nlacolla@iado-conicet.gob.ar).

Comentarios respecto a los Anexos recientemente incorporados del proyecto de resolución “Modificación de la Resolución Presidencia ACUMAR N° 46/2017”

Parte general:

En el artículo 6° del texto de la modificación, se establece sustituir artículo 12 de la Resolución Presidencia ACUMAR N° 46/2017, con el siguiente texto: *“ARTÍCULO 12.- La presente entrará en vigencia al día siguiente de su publicación en el Boletín Oficial de la República Argentina, a excepción de las condiciones de vertido establecidas en el criterio 1 y en el criterio 2 -a los que refiere el artículo 2° de la presente- y la Tabla Consolidada de Control de Límites de Vertido de Efluentes Líquidos -aprobada por ANEXO I (IF xxx) de la presente-, las que entrarán en vigencia el 1 de enero de 2022, con posibilidad de establecer una prórroga como máximo de veinticuatro (24) meses contados a partir de dicha fecha, quedando derogada a partir de dicho momento la Resolución ACUMAR N° 1/2007”.*

Considerando que la recomposición ambiental (objetivo central de ACUMAR) es una necesidad inmediata, se considera que la dilación propuesta es excesiva, y que no hay elementos de peso que la justifiquen adecuadamente, más aun teniendo en cuenta la partición de la cuenca en subcuencas, y que éstas presentan calidades diferentes. Eventualmente puede proponerse un diagrama de aplicación diferenciado para cada una de éstas, pero nunca dilatar la aplicación tres años.

Más aún, consideramos importante volver a plantear nuestro cuestionamiento a las metas de calidad planteadas para la cuenca y en particular los plazos establecidos para su logro, que se establecen en el Artículo 8° de la original Resolución 46/17.

En primer lugar, respecto al Uso IV, “Apta para actividades recreativas pasivas”, no se define con claridad a qué tipo de actividades se refiere, ni es una categoría usada en las legislaciones de otros países. Además, su implementación en el modo que se plantea, permite el deterioro en el corto plazo de la calidad ambiental de la cuenca alta y media, que ya son aptas para este tipo de “uso” y algunos más estrictos. Más elementos se brindan más adelante con los comentarios al Anexo III.

Hay acuerdo con los otros usos, incluso con la mayoría de los valores que se establecen, y de los parámetros que se exigen, más aún con las recientes modificaciones, pero con una sola salvedad: que el cumplimiento de los parámetros deben pedirse en toda la cuenca, más aún con los generales (físicoquímicos y microbiológicos), pero respetando las particularidades de cada subcuenca. Sin embargo, destacamos nuestra discrepancia con los plazos estipulados. Ya se hizo referencia a la aplicación del presunto uso IV, señalándolo como excesiva dilación. Lo mismo ocurre con los otros plazos, resultan excesivamente laxos, más aún cuando por falta de modelización, no se conoce la dinámica de los contaminantes dentro de la cuenca, contando sólo con descripciones de sus concentraciones en agua, y a veces en sedimentos; pero careciendo de

relevamientos topográficos y batimétricos, además de otros datos de la dinámica hidrológica, incluyendo dinámica de sedimentos y especiación de los metales presentes y otras sustancias orgánicas. Desde el mandato de la recomposición ambiental es preferible manejar plazos más estrechos y eventualmente prorrogables, pero acompañando desde ACUMAR con los estudios necesarios para la modelización señalada. Se destaca que desde la red REPBA del CONICET se puede realizar los estudios referidos.

Finalmente, dadas las características de la CMR sostenemos que el grado de contaminación en la cuenca debe estar permanentemente asociado a las dimensiones sociales, por ejemplo tener siempre presente los aspectos asociados al ordenamiento territorial y la salud poblacional.

Respecto al Anexo I

El artículo 3 establece *“Los establecimientos que generen efluentes líquidos que pudieran contener residuos patogénicos, deberán satisfacer la Demanda de Cloro residual de sus efluentes, previo a su descarga”*.

En este caso, se destaca que resultaría de gran utilidad para un mejor contralor del cumplimiento categorizar los tipos de establecimientos, de modo de enfocar con particular atención los parámetros asociados con las características específicas de cada tipo productivo.

Los comentarios restantes, se refieren a la Tabla Consolidada de Control de Límites de Vertido de Efluentes Líquidos.

En primer lugar, y en conexión con el comentario anterior, se considera óptimo y de gran relevancia, establecer límites de vertidos específicos para cada tipo de actividad o industria, con el fin de alcanzar los límites más estrictos que pueden ser viables económicamente para cada una. Este criterio se aplica, por ejemplo, en las normativas de la Agencia Ambiental de los EEUU (US-EPA).

Respecto a límite del **Fósforo total**, se considera que 5 mgP/L para cuerpo de agua superficial es un valor muy elevado, atento a que el medio receptor es un ambiente sujeto a extrema eutrofización y a que se pretende recomponer el sistema. A fines comparativos, la Resolución N° 1089 de la Provincia de Santa Fe (Reglamento para el control del vertimiento de líquidos residuales) establece 2 mgP/L como máximo; mientras que la Resolución N° 336-03 (modificatoria de la N° 389/98) del ADA (Pcia de Bs As) establece $\leq 1,0$ mg/L de P para vertido a cuerpo de agua superficial. Asimismo, la Dirección General del Agua del Ministerio de Medio Ambiente de España establece 1 mgP/L como máximo para vertido a cuerpos superficiales sensibles.

Nuevamente para cuerpo de agua superficial como medio receptor, pero respecto al **Nitrógeno total Kjeldahl**, fijado en ≤ 35 mgN/L, se recomienda establecer límites más estrictos considerando la sensibilidad del ambiente receptor, su estado y la intención de sanearlo. Como comparación se señala que la Dirección General del Agua del Ministerio de Medio Ambiente de España establece 10 mgN/L como máximo para vertido a cuerpos superficiales sensibles. Del mismo modo, la

Resolución 1089 de la Provincia de Santa Fe ya citada, establece que no deben superarse los 15 mgN/L; mientras que el límite superior para la Resolución N° 336-03 (PBA) es de 25 mgN/L.

Finalmente, se recomienda modificar la exigencia de cumplimiento de los límites establecidos para los coliformes fecales (Ref. e), de manera de incluir en ella toda la cuenca y no solamente la cuenca alta, sin perder de vista las características de cada subcuenca. En efecto, si se quiere recomponer la calidad ambiental de la totalidad de la cuenca y alcanzar los niveles de USO 1, 2 o 3, los coliformes fecales deben regularse en la totalidad de la cuenca.

Respecto al Anexo III

Comienza con una afirmación muy extensa y poco precisa. Dice: *“En todos los cuerpos de agua superficial y para todos los Usos establecidos en la presente, deberá verificarse la ausencia de los siguientes componentes: materia flotante y espumas no naturales; aceites minerales, vegetales, y grasas; colorantes de fuentes antrópicas; y residuos sólidos de fuentes antrópicas.*

Entiéndase por el término “Ausencia” que los componentes indicados en el párrafo precedente no deben estar presentes en concentraciones que sean detectables a simple vista o por olor; o bien que puedan formar depósitos en las orillas de los arroyos o ríos; o bien afectar a los organismos acuáticos presentes.

Es poco preciso el término ausencia, y queda atado a la percepción de cada quien, en términos subjetivos y cualitativos. En todo caso, es necesario definir esta disposición en términos cuantitativos, una metodología para su detección y una escala de valores. Por ejemplo, en términos perceptivos, como se reconoce si es una espuma natural o no natural? Como se reconoce si un colorante es de fuente antrópica?.

Finalmente, la propuesta de incorporar los estudios ecotoxicológicos, es una forma concreta de dar respuesta cuantitativa al requisito *“... o bien afectar a los organismos acuáticos presentes”*.

Por otro lado, y respecto a los distintos Usos del agua, se considera que los parámetros de análisis deben contar con límites cuantificados para la totalidad de los usos (del I al IV), dado que se debe recomponer la calidad ambiental de la totalidad de la cuenca, sin perder de vista las características de cada subcuenca, como ya se señaló anteriormente para el parámetro coliformes totales, y como ocurre con el parámetros SSEE (aceites y grasas).

En particular, se cuestiona el uso IV “Apta para actividades recreativas pasivas” dado que no se define con claridad los alcances de las mencionadas “actividades recreativas pasivas”, y que no es una categoría usada en las legislaciones de otros países. Es más, es un objetivo muy poco ambicioso, que permite el deterioro, en el corto plazo, de la calidad ambiental de la cuenca alta y media, que ya son aptas para este tipo de “uso” y algunos más estrictos. Por otra parte, en la mayor parte de la cuenca baja se ha construido un camino de sirga donde se desarrollan actividades recreativas o productivas sin contacto directo con el río (Uso III) por lo que este artículo no tendría aplicación concreta.

Propuesta de incorporación de estudios ecotoxicológicos en el ANEXO 1:

Antecedentes y fundamento

El saneamiento de la cuenca del Río Matanza-Riachuelo tiene como objetivo ecológico recuperar condiciones de vida compatibles con la integridad del ecosistema. Siendo la toxicidad en el agua y sedimentos uno de los condicionantes fundamentales que impiden la recuperación de la biota y el consiguiente saneamiento, es fundamental incorporar el Riesgo por Toxicidad como un criterio de gestión prioritario para alcanzar el objetivo de evitar el vuelco de efluentes y otros materiales en concentraciones tóxicas, incompatibles con el ecosistema acuático. Los estudios ecotoxicológicos abarcan un amplísimo rango de variables que incluyen, diferentes especies, diferentes condiciones, y numerosísimos objetivos específicos, etc. En América Latina y en Argentina en particular, ya en 1997 había suficientes grupos de científicos interesados en Toxicología y Química Ambiental como para formar parte de una sociedad internacional, SETAC, organizando cursos, simposios, congresos, etc. de tal manera que a la fecha hay centenares de contribuciones científicas en Ecotoxicología desde distintos Centros de Investigación en nuestro país, lo que implica una importante infraestructura de laboratorios y personal entrenado en esta disciplina (ANEXO A).

En 1991 la EPA ha publicado un documento denominado "Technical Support Document for Water Quality-based Toxics Control" que permite, en base a una **metodología claramente definida**, establecer límites de toxicidad admisible para los permisos de vuelco a fin que los mismos sean compatibles con el ecosistema (1). Expresado en Unidades de Toxicidad, el Riesgo por Toxicidad Aguda (96 hs. de exposición) no debe superar el valor de 0,3 y para crónica corta (7 días de exposición) el valor de 1. Aplicando dicho criterio, a partir de 1996 se han publicado trabajos desde la Argentina informando el Riesgo por Toxicidad en efluentes industriales, lixiviados aguas y sedimentos de cuencas hídricas de sitios considerados no impactados hasta localizaciones severamente afectadas por la actividad antrópica (Anexo A). La información es científicamente robusta, económica para obtener, reproducible, permite gestionar objetivos cada vez más ambiciosos en el espacio y el tiempo para alcanzar la sustentabilidad en toda la cuenca. Es fácil de entender para todas las partes interesadas, organismos de gestión y control, empresas con sus diferentes roles y la población asentada en la cuenca. No es menos relevante que los esfuerzos para alcanzar el saneamiento tenga definido uno de los objetivos centrales, que el riesgo por toxicidad sea compatible con la vida, orientando las acciones para alcanzar dicho objetivo. La recuperación de la biota es fundamental para todos los mecanismos de resiliencia que participaran activamente en el saneamiento de la cuenca. La incorporación de este criterio de gestión es complementaria con cualquier otro que se estuviera ejecutando o a incorporar.

1. Technical Support Document for Water Quality-based Toxics Control. EPA /505/2-90-001, 1991.

Texto a incorporar en el ANEXO 1 de la Res. ACUMAR 46/2017

Artículo 8. Incorpórese la realización de estudios Ecotoxicológicos en el agua, sedimentos, efluentes industriales y lixiviados que impactan la cuenca del Rio Matanza-Riachuelo. A fin de establecer la línea de base para integrar a la gestión de la cuenca el criterio de RIESGO por Toxicidad, se realizara durante un periodo de 2 años un monitoreo de la Toxicidad en las 14 subcuencas en el agua, sedimentos, efluentes industriales y lixiviados que impacten el ecosistema. La metodología y limites por toxicidad a ser implementada se basa en el "Technical Support Document for Water Quality-based Toxics Control" de la US EPA 1991. El Riesgo por Toxicidad aguda (96 hs de exposición), expresado en Unidades de Toxicidad ($UTa=100/CL50$) tendría como valor máximo admisible 0,3 y para toxicidad crónica corta (7 días de exposición), expresado en Unidades de Toxicidad ($UTC=100/NOEC$) 1. Los objetivos a futuro para agua y sedimentos podrán ser perfeccionados ampliando el tiempo de exposición a crónico (10 o más días de exposición) y a efectos subletales, v.g. teratogénesis/ otros biomarcadores de toxicidad.

ANEXO A

Bibliografía que respalda la experiencia en la Argentina sobre la evaluación de efectos biológicos de agentes físicos y químicos solos o en combinación, desarrollo e implementación de bioensayos en laboratorio y/o campo, su aplicación para la evaluación de calidad de agua, sedimentos, efluentes industriales, lixiviados, diagnóstico ambiental, etc. Las publicaciones marcadas con asterisco corresponden a contribuciones en las que se informó toxicidad aplicando los criterios del documento de la EPA "Technical Support Document for Water Quality-based Toxics Control (1991).

Achiorno, C. L., De Villalobos, C., & Ferrari, L. (2016). Analysis Of The Infectivity Of Chordodes Nobilii Larvae (Gordiida, Nematomorpha). Revista de Biología Tropical, 65(1), 1. <https://doi.org/10.15517/rbt.v65i1.23536>

Achiorno, C., Ferrari, L., & Villalobos, C. (2008). Effect of extreme temperature on egg development, larval and adult survival of Chordodes nobilii Camerano, 1901 (Gordiida, Nematomorpha). Acta Parasitologica, 53(4). <https://doi.org/10.2478/s11686-008-0052-5>

Agostini, M. G., Natale, G. S., & Ronco, A. E. (2009). Impact of endosulphan and cypermethrin mixture on amphibians under field use for biotech soya bean production. International Journal of Environment and Health, 3(4), 379. <https://doi.org/10.1504/IJENVH.2009.030109>

Agostini, M. G., Natale, G. S., & Ronco, A. E. (2010). Lethal and sublethal effects of cypermethrin to Hypsiboas pulchellus tadpoles. Ecotoxicology, 19(8), 1545–1550. <https://doi.org/10.1007/s10646-010-0539-3>

Agostini, M., Kacoliris, F., Demetrio, P., Natale, G., Bonetto, C., & Ronco, A. (2013). Abnormalities in amphibian populations inhabiting agroecosystems in northeastern Buenos Aires Province, Argentina. Diseases of Aquatic Organisms, 104(2), 163–171. <https://doi.org/10.3354/dao02592>

Albariño, R., Venturino, A., Montagna, C. M., & Pechen de D'Angelo, A. M. (2007). Environmental effect assessment of magnacide® herbicide at río colorado irrigation channels (argentina). Tier 4: in situ survey on benthic invertebrates. Environmental Toxicology and Chemistry, 26(1), 183. <https://doi.org/10.1897/06-086R.1>

Alberto, W. D., María del Pilar, D., María Valeria, A., Fabiana, P. S., Cecilia, H. A., & María de los Ángeles, B. (2001). Pattern Recognition Techniques for the Evaluation of

Spatial and Temporal Variations in Water Quality. A Case Study: Water Research, 35(12), 2881–2894. [https://doi.org/10.1016/S0043-1354\(00\)00592-3](https://doi.org/10.1016/S0043-1354(00)00592-3)

- Ale, A., Bacchetta, C., Rossi, A. S., Galdopórpora, J., Desimone, M. F., de la Torre, F. R., ... Cazenave, J. (2018). Nanosilver toxicity in gills of a neotropical fish: Metal accumulation, oxidative stress, histopathology and other physiological effects. *Ecotoxicology and Environmental Safety*, 148, 976–984. <https://doi.org/10.1016/j.ecoenv.2017.11.072>
- Ale, A., Liberatori, G., Vannuccini, M. L., Bergami, E., Ancora, S., Mariotti, G., ... Corsi, I. (2019). Exposure to a nanosilver-enabled consumer product results in similar accumulation and toxicity of silver nanoparticles in the marine mussel *Mytilus galloprovincialis*. *Aquatic Toxicology*, 211, 46–56. <https://doi.org/10.1016/j.aquatox.2019.03.018>
- Ale, A., Rossi, A. S., Bacchetta, C., Gervasio, S., de la Torre, F. R., & Cazenave, J. (2018). Integrative assessment of silver nanoparticles toxicity in *Prochilodus lineatus* fish. *Ecological Indicators*, 93, 1190–1198. <https://doi.org/10.1016/j.ecolind.2018.06.023>
- Amé, M. V., & Pesce, S. F. (2015). Spatial and Temporal Changes in Water Quality Along the Basin (pp. 93–111). https://doi.org/10.1007/698_2015_434
- Amé, M. V., & Wunderlin, D. A. (2005). Effects of Iron, Ammonium and Temperature on Microcystin Content by a Natural Concentrated *Microcystis Aeruginosa* Population. *Water, Air, and Soil Pollution*, 168(1–4), 235–248. <https://doi.org/10.1007/s11270-005-1774-8>
- Amé, M. V., Baroni, M. V., Galanti, L. N., Bocco, J. L., & Wunderlin, D. A. (2009). Effects of microcystin–LR on the expression of P-glycoprotein in *Jenynsia multidentata*. *Chemosphere*, 74(9), 1179–1186. <https://doi.org/10.1016/j.chemosphere.2008.11.068>
- Amé, M. V., del Pilar Díaz, M., & Wunderlin, D. A. (2003). Occurrence of toxic cyanobacterial blooms in San Roque Reservoir (Córdoba, Argentina): A field and chemometric study. *Environmental Toxicology*, 18(3), 192–201. <https://doi.org/10.1002/tox.10114>
- Amé, M. V., Galanti, L. N., Menone, M. L., Gerpe, M. S., Moreno, V. J., & Wunderlin, D. A. (2010). Microcystin–LR, –RR, –YR and –LA in water samples and fishes from a shallow lake in Argentina. *Harmful Algae*, 9(1), 66–73. <https://doi.org/10.1016/j.hal.2009.08.001>

- Andrade, V. S., Gutierrez, M. F., Fantón, N. I., & Gagneten, A. M. (2018). Shifts in Zooplankton Behavior Caused by a Mixture of Pesticides. *Water, Air, & Soil Pollution*, 229(4), 107. <https://doi.org/10.1007/s11270-018-3752-y>
- Anguiano, O. L., Castro, C., Venturino, A., & Ferrari, A. (2014). Acute toxicity and biochemical effects of azinphos methyl in the amphipod *Hyaella curvispina*. *Environmental Toxicology*, 29(9), 1043–1053. <https://doi.org/10.1002/tox.21834>
- Anguiano, O. L., Ferrari, A., Soleño, J., Martínez, M. C., Venturino, A., Pechen de D'Angelo, A. M., & Montagna, C. M. (2008). Enhanced Esterase Activity And Resistance To Azinphosmethyl In Target And Nontarget Organisms. *Environmental Toxicology and Chemistry*, 27(10), 2117. <https://doi.org/10.1897/07-655.1>
- Anguiano, O. L., Vacca, M., Rodriguez Araujo, M. E., Montagna, M., Venturino, A., & Ferrari, A. (2017). Acute toxicity and esterase response to carbaryl exposure in two different populations of amphipods *Hyaella curvispina*. *Aquatic Toxicology*, 188, 72–79. <https://doi.org/10.1016/j.aquatox.2017.04.013>
- Aronzon C M, Babay P A, Pérez Coll CS Developmental Toxicity and Risk Assessment of Nonylphenol to the South American Toad, *Rhinella arenarum*. *Environm. Toxicol Pharmacol.* 24:38(2) 634-642, 2014. DOI: 10.1016/j.etap.2014.08.014
- Aronzon C. M, Svartz G., Pérez Coll C. S. “Joint toxicity of copper and diazinon during embryonic and larval development of the common South American toad, *Rhinella arenarum*” *Chemistry and Ecology*. 32 (9): 829-842 Issue 9, 2016
- Aronzon C.; Marino D.J.G; Ronco A.E.; Pérez Coll C.S. Differential toxicity and uptake of Diazinon on embryo-larval development of *Rhinella arenarum* *Chemosphere* 100:50-56, 2014.
- Aronzon CM, Sandoval MT, Herkovits J, Pérez-Coll CS. Stage-dependent toxicity of 2,4-dichlorophenoxyacetic on the embryonic development of a South American toad, *Rhinella arenarum*. *Environ Toxicol.* 26:n/a. doi: 10.1002/tox.20564 2010.
- Aronzon CM, Svartz GV, Pérez Coll, CS Synergy Between Diazinon and Nonylphenol in Toxicity During the Early Development of the *Rhinella arenarum* Toad. *Water, Air, & Soil Pollution*, 227(5), 1-10 DOI 10.1007/s11270-016-2799-x <http://link.springer.com/article/10.1007/s11270-016-2799-x> 227:139 ISSN: 1573-2932
- Aronzon CM, Sandoval MT, Herkovits J*, Pérezcoll CS.(*corresponding author). Stage-dependent susceptibility to copper in *Rhinella arenarum* embryos and larvae. *Environ Toxicol Chem.* , 30, 12, 2771-7. 2011

- Aronzon, C., Svartz, G., Pérez Coll, CS. Comparative toxicity of endosulfan and diazinon on the embryo-larval development of the South American toad, *Rhinella arenarum*. *Int. J. Environ. Health* 8(3):225-234, 2017
- Arreghini, S., de Cabo, L., de Iorio, A. F., Rendina, A., & Bonetto, C. (2001). Effects of Zinc on the Growth of Bulrush (*Schoenoplectus californicus*) and Its Distribution Between Different Sediment Fractions. *Bulletin of Environmental Contamination and Toxicology*, 67(2), 264–270. <https://doi.org/10.1007/s001280119>
- Asorey, C.M., Muñoz, L.M., Pérez-Coll, CS., Lipscomb, J. y Herkovits, J. Efecto del ácido dicloroacético sobre la toxicidad del cobre y el arsénico en embriones de anfibio. En “Salud Ambiental y Humana: Una visión holística” pp.94-96. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1
- Bacchetta, C., Ale, A., Simoniello, M. F., Gervasio, S., Davico, C., Rossi, A. S., ... Cazenave, J. (2017). Genotoxicity and oxidative stress in fish after a short-term exposure to silver nanoparticles. *Ecological Indicators*, 76, 230–239. <https://doi.org/10.1016/j.ecolind.2017.01.018>
- Bacchetta, C., Cazenave, J., & Parma, M. J. (2011). Responses of Biochemical Markers in the Fish *Prochilodus lineatus* Exposed to a Commercial Formulation of Endosulfan. *Water, Air, & Soil Pollution*, 216(1–4), 39–49. <https://doi.org/10.1007/s11270-010-0512-z>
- Bacchetta, C., Cazenave, J., Parma, M. J., & Biancucci, G. F. (2011). Biochemical Stress Responses in Tissues of the Cichlid Fish *Cichlasoma dimerus* Exposed to a Commercial Formulation of Endosulfan. *Archives of Environmental Contamination and Toxicology*, 61(3), 453–460. <https://doi.org/10.1007/s00244-010-9635-x>
- Bacchetta, C., Rossi, A. S., Cian, R. E., Drago, S. R., & Cazenave, J. (2019). Dietary β -carotene improves growth performance and antioxidant status of juvenile *Piaractus mesopotamicus*. *Aquaculture Nutrition*, anu.12893. <https://doi.org/10.1111/anu.12893>
- Bacchetta, C., Rossi, A., Ale, A., Campana, M., Parma, M. J., & Cazenave, J. (2014). Combined toxicological effects of pesticides: A fish multi-biomarker approach. *Ecological Indicators*, 36, 532–538. <https://doi.org/10.1016/j.ecolind.2013.09.016>
- Bach, N. C., Marino, D. J. G., Natale, G. S., & Somoza, G. M. (2018). Effects of glyphosate and its commercial formulation, Roundup® Ultramax, on liver histology of tadpoles of the neotropical frog, *Leptodactylus latrans* (amphibia: Anura). *Chemosphere*, 202, 289–297. <https://doi.org/10.1016/j.chemosphere.2018.03.110>

- Bach, N. C., Natale, G. S., Somoza, G. M., & Ronco, A. E. (2016). Effect on the growth and development and induction of abnormalities by a glyphosate commercial formulation and its active ingredient during two developmental stages of the South-American Creole frog, *Leptodactylus latrans*. *Environmental Science and Pollution Research*, 23(23), 23959–23971. <https://doi.org/10.1007/s11356-016-7631-z>
- Baigún, C., Mugni, H., & Bonetto, C. (2006). Nutrient Concentrations and Trophic State of Small Patagonian Andean lakes. *Journal of Freshwater Ecology*, 21(3), 449–456. <https://doi.org/10.1080/02705060.2006.9665022>
- Ballesteros, M. L., Gonzalez, M., Wunderlin, D. A., Bistoni, M. A., & Miglioranza, K. S. B. (2011). Uptake, tissue distribution and metabolism of the insecticide endosulfan in *Jenynsia multidentata* (Anablepidae, Cyprinodontiformes). *Environmental Pollution*, 159(6), 1709–1714. <https://doi.org/10.1016/j.envpol.2011.02.037>
- Ballesteros, M. L., Hued, A. C., Gonzalez, M., Miglioranza, K. S. B., & Bistoni, M. A. (2017). Evaluation of the Health Status of the Silverside (*Odontesthes bonariensis*) at a RAMSAR Site in South America. *Bulletin of Environmental Contamination and Toxicology*, 99(1), 62–68. <https://doi.org/10.1007/s00128-017-2055-0>
- Ballesteros, M. L., Rivetti, N. G., Morillo, D. O., Bertrand, L., Amé, M. V., & Bistoni, M. A. (2017). Multi-biomarker responses in fish (*Jenynsia multidentata*) to assess the impact of pollution in rivers with mixtures of environmental contaminants. *Science of The Total Environment*, 595, 711–722. <https://doi.org/10.1016/j.scitotenv.2017.03.203>
- Barni, M. F. S., Ondarza, P. M., Gonzalez, M., Da Cuña, R., Meijide, F., Grosman, F., ... Miglioranza, K. S. B. (2016). Persistent organic pollutants (POPs) in fish with different feeding habits inhabiting a shallow lake ecosystem. *Science of The Total Environment*, 550, 900–909. <https://doi.org/10.1016/j.scitotenv.2016.01.176>
- Baudou, F. G., Ossana, N. A., Castañé, P. M., Mastrángelo, M. M., & Ferrari, L. (2017). Cadmium effects on some energy metabolism variables in *Cnesterodon decemmaculatus* adults. *Ecotoxicology*, 26(9), 1250–1258. <https://doi.org/10.1007/s10646-017-1850-z>
- Baudou, F. G., Ossana, N. A., Castañé, P. M., Mastrángelo, M. M., González Núñez, A. A., Palacio, M. J., & Ferrari, L. (2019). Use of integrated biomarker indexes for assessing the impact of receiving waters on a native neotropical teleost fish. *Science of The Total Environment*, 650, 1779–1786. <https://doi.org/10.1016/j.scitotenv.2018.09.342>
- Belmonte, N.; Rivera, O. and and Herkovits, J. Zinc Protection against Cd effects on preimplantation mice embryos.. *Bull. Environ. Contam.Toxicol.* 43: 107 - 110, 1989.

- Bertrand, L., Asis, R., Monferrán, M. V., & Amé, M. V. (2016). Bioaccumulation and biochemical response in South American native species exposed to zinc: Boosted regression trees as novel tool for biomarkers selection. *Ecological Indicators*, 67, 769–778. <https://doi.org/10.1016/j.ecolind.2016.03.048>
- Bertrand, L., Marino, D. J., Monferrán, M. V., & Amé, M. V. (2017). Can a low concentration of an organophosphate insecticide cause negative effects on an aquatic macrophyte? Exposure of *Potamogeton pusillus* at environmentally relevant chlorpyrifos concentrations. *Environmental and Experimental Botany*, 138, 139–147. <https://doi.org/10.1016/j.envexpbot.2017.03.006>
- Bertrand, L., Monferrán, M. V., Métais, I., Mouneyrac, C., & Amé, M. V. (2015). MTs in *Palaemonetes argentinus* as potential biomarkers of zinc contamination in freshwaters. *Ecological Indicators*, 48, 533–541. <https://doi.org/10.1016/j.ecolind.2014.09.019>
- Bertrand, L., Monferrán, M. V., Mouneyrac, C., & Amé, M. V. (2018). Native crustacean species as a bioindicator of freshwater ecosystem pollution: A multivariate and integrative study of multi-biomarker response in active river monitoring. *Chemosphere*, 206, 265–277. <https://doi.org/10.1016/j.chemosphere.2018.05.002>
- Bertrand, L., Monferrán, M. V., Mouneyrac, C., Bonansea, R. I., Asis, R., & Amé, M. V. (2016). Sensitive biomarker responses of the shrimp *Palaemonetes argentinus* exposed to chlorpyrifos at environmental concentrations: Roles of alpha-tocopherol and metallothioneins. *Aquatic Toxicology*, 179, 72–81. <https://doi.org/10.1016/j.aquatox.2016.08.014>
- Bertrand, L., Monferrán, M. V., Valdés, M. E., & Amé, M. V. (2019). Usefulness of a freshwater macrophyte (*Potamogeton pusillus*) for an environmental risk assessment in a multi-source contaminated basin. *Chemosphere*, 222, 1003–1016. <https://doi.org/10.1016/j.chemosphere.2019.02.018>
- Birba, A., Ramallo, M. R., Lo Nostro, F., Guimarães Moreira, R., & Pandolfi, M. (2015). Reproductive and parental care physiology of *Cichlasoma dimerus* males. *General and Comparative Endocrinology*, 221, 193–200. <https://doi.org/10.1016/j.ygcen.2015.02.004>
- Bonansea, R. I., Marino, D. J. G., Bertrand, L., Wunderlin, D. A., & Amé, M. V. (2017). Tissue-specific bioconcentration and biotransformation of cypermethrin and chlorpyrifos in a native fish (*Jenynsia multidentata*) exposed to these insecticides singly and in mixtures. *Environmental Toxicology and Chemistry*, 36(7), 1764–1774. <https://doi.org/10.1002/etc.3613>

- Bonansea, R. I., Wunderlin, D. A., & Amé, M. V. (2016). Behavioral swimming effects and acetylcholinesterase activity changes in *Jenynsia multidentata* exposed to chlorpyrifos and cypermethrin individually and in mixtures. *Ecotoxicology and Environmental Safety*, 129, 311–319. <https://doi.org/10.1016/j.ecoenv.2016.03.043>
- Bonifacio, A. F., Ballesteros, M. L., Bonansea, R. I., Filippi, I., Amé, M. V., & Hued, A. C. (2017). Environmental relevant concentrations of a chlorpyrifos commercial formulation affect two neotropical fish species, *Cheirodon interruptus* and *Cnesterodon decemmaculatus*. *Chemosphere*, 188, 486–493. <https://doi.org/10.1016/j.chemosphere.2017.08.156>
- Bonifacio, A. F., Cazenave, J., Bacchetta, C., Ballesteros, M. L., de los Ángeles Bistoni, M., Amé, M. V., ... Hued, A. C. (2016). Alterations in the general condition, biochemical parameters and locomotor activity in *Cnesterodon decemmaculatus* exposed to commercial formulations of chlorpyrifos, glyphosate and their mixtures. *Ecological Indicators*, 67, 88–97. <https://doi.org/10.1016/j.ecolind.2016.02.011>
- Bonifacio, A. F., Cazenave, J., Bacchetta, C., Ballesteros, M. L., de los Ángeles Bistoni, M., Amé, M. V., ... Hued, A. C. (2016). Alterations in the general condition, biochemical parameters and locomotor activity in *Cnesterodon decemmaculatus* exposed to commercial formulations of chlorpyrifos, glyphosate and their mixtures. *Ecological Indicators*, 67, 88–97. <https://doi.org/10.1016/j.ecolind.2016.02.011>
- Brodeur, J C Asorey C M, Sztrum A, and Herkovits J. Acute and Subchronic Toxicity of Arsenite and Zinc to Tadpoles of *Rhinella arenarum* Both Alone and in Combination. *Journal of Toxicology and Environmental Health, Part A*, 72: 884–890, 2009 ISSN: 1528-7394 print / 1087-2620 online
- Brodeur, J. C., Suarez, R. P., Natale, G. S., Ronco, A. E., & Elena Zaccagnini, M. (2011). Reduced body condition and enzymatic alterations in frogs inhabiting intensive crop production areas. *Ecotoxicology and Environmental Safety*, 74(5), 1370–1380. <https://doi.org/10.1016/j.ecoenv.2011.04.024>
- Brodeur, J.C., Svart, G; Perez-Coll, C.S., Marino, D.J.G., Herkovits, J. Comparative susceptibility to atrazine of three developmental stages of *Bufo arenarum* (amphibia: anura) and influence on metamorphosis: non-monotonous acceleration of the time to climax and delayed tail resorption *Aquat. Toxicol.*, 91(2):161-170, 2009.
- Bustoabad, O.; Herkovits, J. and Pisanó, A. Different sensitivity to lithium ion during the segmentation of *Bufo arenarum* eggs. *Acta Embryol. Exp.*, 3: 271 - 282, 1977.

- C. Pérez-Coll, J. Herkovits y O. Domínguez. Efecto sinérgico entre el 2,4-D y el cobre sobre embriones de anfibio., *Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable*. SETAC Publication, Ed. Jorge Herkovits, 5-7, 2003
- C..S. , Herkovits, J.; Fridman, O.; Daniel, P. and D'Eramo J.L. Metallothionein induction and cadmium uptake in *Bufo arenarum* embryos following an acclimation protocol. *Environmental Pollution* 106, 443-448, 1999.
- Carabajal Paladino, L., Lo Nostro, F., Papeschi, A., Cladera, J., & Bressa, M. J. (2017). Spermatogenesis in the parasitoid *Diachasmimorpha longicaudata* (Hymenoptera: Braconidae). *Acta Zoologica*, 98(1), 38–43. <https://doi.org/10.1111/azo.12148>
- Carriquiriborde, P., Díaz, J., Mugni, H., Bonetto, C., & Ronco, A. E. (2007). Impact of cypermethrin on stream fish populations under field-use in biotech-soybean production. *Chemosphere*, 68(4), 613–621. <https://doi.org/10.1016/j.chemosphere.2007.02.051>
- Castañaga L., Cordon G., Corin J., Pérez Coll C. S., Sztrum A. y Herkovits J. 2009. Efecto letal y teratogénico del H₂O₂ sobre el desarrollo embrionario de *Chaunus arenarum* (Anura, Bufonidae). *Química y Toxicología Ambiental en América Latina*, Ed. J.Herkovits, pp 89-91. ISBN 978-987-25370-0-5.
- Castañaga, Luis A., Asorey, C., Sandoval, María T., Pérez-Coll, C. S., Argibay, T. and Herkovits, J. Stage Dependent Teratogenic and Lethal Effects Exerted By Ultraviolet B Radiation on *Rhinella* (*Bufo*) *arenarum* Embryos. *Environ. Toxicol. Chem.* 28(2):427-433, 2009.
- Castillo Morales, Gabriela (ed.) *Ensayos toxicológicos y métodos de evaluación de calidad de aguas. Estandarización, intercalibración, resultados y aplicaciones*/Gabriela Castillo (ed.). - México: IMTA, 2004.
- Castro, J. M., Bianchi, V. A., Pascual, M. M., Almeida, C., Venturino, A., & Luquet, C. M. (2018). Immune and biochemical responses in hemolymph and gills of the Patagonian freshwater mussel *Diplodon chilensis*, against two microbiological challenges: *Saccharomyces cerevisiae* and *Escherichia coli*. *Journal of Invertebrate Pathology*, 157, 36–44. <https://doi.org/10.1016/j.jip.2018.08.005>
- Castro, J. M., Bianchi, V. A., Pascual, M., Venturino, A., & Luquet, C. M. (2017). Modulation of immune and antioxidant responses by azinphos-methyl in the freshwater mussel *Diplodon chilensis* challenged with *Escherichia coli*. *Environmental Toxicology and Chemistry*, 36(7), 1785–1794. <https://doi.org/10.1002/etc.3612>

- Cazenave, J., Bacchetta, C., Parma, M. J., Scarabotti, P. A., & Wunderlin, D. A. (2009). Multiple biomarkers responses in *Prochilodus lineatus* allowed assessing changes in the water quality of Salado River basin (Santa Fe, Argentina). *Environmental Pollution*, 157(11), 3025–3033. <https://doi.org/10.1016/j.envpol.2009.05.055>
- Cazenave, J., Bacchetta, C., Rossi, A., Ale, A., Campana, M., & Parma, M. J. (2014). Deleterious effects of wastewater on the health status of fish: A field caging study. *Ecological Indicators*, 38, 104–112. <https://doi.org/10.1016/j.ecolind.2013.10.029>
- Cazenave, J., Bistoni, M. de los A., Pesce, S. F., & Wunderlin, D. A. (2006). Differential detoxification and antioxidant response in diverse organs of *Corydoras paleatus* experimentally exposed to microcystin-RR. *Aquatic Toxicology*, 76(1), 1–12. <https://doi.org/10.1016/j.aquatox.2005.08.011>
- Cazenave, J., Bistoni, M. de los Á., Zwirnmann, E., Wunderlin, D. A., & Wiegand, C. (2006). Attenuating effects of natural organic matter on microcystin toxicity in zebra fish (*Danio rerio*) embryos—benefits and costs of microcystin detoxication. *Environmental Toxicology*, 21(1), 22–32. <https://doi.org/10.1002/tox.20151>
- Cazenave, J., Nores, M. L., Miceli, M., Díaz, M. P., Wunderlin, D. A., & Bistoni, M. A. (2008). Changes in the swimming activity and the glutathione S-transferase activity of *Jenynsia multidentata* fed with microcystin-RR. *Water Research*, 42(4–5), 1299–1307. <https://doi.org/10.1016/j.watres.2007.09.025>
- Cazenave, J., Wunderlin, D. A., Bistoni, M. de los Á., Amé, M. V., Krause, E., Pflugmacher, S., & Wiegand, C. (2005). Uptake, tissue distribution and accumulation of microcystin-RR in *Corydoras paleatus*, *Jenynsia multidentata* and *Odontesthes bonariensis*. *Aquatic Toxicology*, 75(2), 178–190. <https://doi.org/10.1016/j.aquatox.2005.08.002>
- Cazenave, J., Wunderlin, D. A., Bistoni, M. de los Á., Amé, M. V., Krause, E., Pflugmacher, S., & Wiegand, C. (2006). Erratum to “Uptake, tissue distribution and accumulation of microcystin-RR in *Corydoras paleatus*, *Jenynsia multidentata* and *Odontesthes bonariensis*.” *Aquatic Toxicology*, 77(4), 439. <https://doi.org/10.1016/j.aquatox.2006.02.005>
- Cazenave, J., Wunderlin, D. A., Hued, A. C., & Bistoni, M. de los A. (2005). Haematological parameters in a neotropical fish, *Corydoras paleatus* (Jenyns, 1842) (Pisces, Callichthyidae), captured from pristine and polluted water. *Hydrobiologia*, 537(1–3), 25–33. <https://doi.org/10.1007/s10750-004-1638-z>
- Chernoff, N., Hunter, E. S., Hall, L.L., Rosen, M.B., Brownie, C. F., Malarkey, D., Marr, M. and Herkovits, J. Lack of Teratogenicity of Microcystin-LR in the Mose and Toad. *Journal of Applied Toxicology*, 22, 13-17, 2002

- Contardo-Jara, V., Galanti, L. N., Amé, M. V., Monferrán, M. V., Wunderlin, D. A., & Wiegand, C. (2009). Biotransformation and antioxidant enzymes of *Limnoperna fortunei* detect site impact in watercourses of Córdoba, Argentina. *Ecotoxicology and Environmental Safety*, 72(7), 1871–1880. <https://doi.org/10.1016/j.ecoenv.2009.07.001>
- Corró, J. Herkovits, O. Fridman. Toxicidad y Captación de estradiol en embriones de *Bufo arenarum*. *Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable*. SETAC Publication, Ed. Jorge Herkovits, 121-123, 2003
- Cristina S. Pérez-Coll, Abelardo A. Sztrum, Jorge Herkovits. Nickel tissue residue as a biomarker of sub-toxic exposure and susceptibility in amphibian embryos *Chemosphere*, 71, 1, 78-83, 2008.
- Crupkin, A. C., Carriquiriborde, P., Mendieta, J., Panzeri, A. M., Ballesteros, M. L., Miglioranza, K. S. B., & Menone, M. L. (2013). Oxidative stress and genotoxicity in the South American cichlid, *Australoheros facetus*, after short-term sublethal exposure to endosulfan. *Pesticide Biochemistry and Physiology*, 105(2), 102–110. <https://doi.org/10.1016/j.pestbp.2012.12.005>
- Crupkin, A. C., Iturburu, F. G., Crupkin, M., & Menone, M. L. (2018). Myofibrillar functional dysregulation in fish: A new biomarker of damage to pesticides. *Ecotoxicology and Environmental Safety*, 158, 44–49. <https://doi.org/10.1016/j.ecoenv.2018.03.096>
- D'Eramo, J. L. Fridman O., Herkovits J. Protección Sinérgica del Zinc y el selenio ante los efectos letales producidos por la radiación UV-B en embriones de *Bufo arenarum*. En "Salud Ambiental y Humana: Una visión holística" P. 91-93 Publicación SETAC. Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1
- D'Eramo, J. L., Fridman, O., Herkovits, J. Respuesta Antioxidante en embriones de *Bufo arenarum* expuestos a dosis letales de radiación con luz ultravioleta (UV-b). *Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable*. SETAC Publication, Ed. Jorge Herkovits, 147-149, 2003
- Da Cuña, R. H., Pandolfi, M., Genovese, G., Piazza, Y., Ansaldo, M., & Lo Nostro, F. L. (2013). Endocrine disruptive potential of endosulfan on the reproductive axis of *Cichlasoma dimerus* (Perciformes, Cichlidae). *Aquatic Toxicology*, 126, 299–305. <https://doi.org/10.1016/j.aquatox.2012.09.015>
- Da Cuña, R. H., Rey Vázquez, G., Dorelle, L., Rodríguez, E. M., Guimarães Moreira, R., & Lo Nostro, F. L. (2016). Mechanism of action of endosulfan as disruptor of gonadal steroidogenesis in the cichlid fish *Cichlasoma dimerus*. *Comparative Biochemistry and*

Physiology Part C: Toxicology & Pharmacology, 187, 74–80.
<https://doi.org/10.1016/j.cbpc.2016.05.008>

Da Cuña, R. H., Rey Vázquez, G., Piol, M. N., Guerrero, N. V., Maggese, M. C., & Lo Nostro, F. L. (2011). Assessment of the acute toxicity of the organochlorine pesticide endosulfan in *Cichlasoma dimerus* (Teleostei, Perciformes). *Ecotoxicology and Environmental Safety*, 74(4), 1065–1073.
<https://doi.org/10.1016/j.ecoenv.2011.02.002>

De Anna, J. S., Leggieri, L. R., Arias Darraz, L., Cárcamo, J. G., Venturino, A., & Luquet, C. M. (2018). Effects of sequential exposure to water accommodated fraction of crude oil and chlorpyrifos on molecular and biochemical biomarkers in rainbow trout. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 212, 47–55. <https://doi.org/10.1016/j.cbpc.2018.07.003>

de Campos, M. F., Lo Nostro, F. L., Da Cuña, R. H., & Moreira, R. G. (2018). Endocrine disruption of phenanthrene in the protogynous dusky grouper *Epinephelus marginatus* (Serranidae: Perciformes). *General and Comparative Endocrinology*, 257, 255–263. <https://doi.org/10.1016/j.ygcen.2017.06.020>

de la Torre, F. ., Salibián, A., & Ferrari, L. (2000). Biomarkers assessment in juvenile *Cyprinus carpio* exposed to waterborne cadmium. *Environmental Pollution*, 109(2), 277–282. [https://doi.org/10.1016/S0269-7491\(99\)00263-8](https://doi.org/10.1016/S0269-7491(99)00263-8)

de la Torre, F. R., Ferrari, L., & Salibián, A. (2002). Freshwater pollution biomarker: response of brain acetylcholinesterase activity in two fish species. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 131(3), 271–280. [https://doi.org/10.1016/S1532-0456\(02\)00014-5](https://doi.org/10.1016/S1532-0456(02)00014-5)

de la Torre, F. R., Ferrari, L., & Salibián, A. (2005). Biomarkers of a native fish species (*Cnesterodon decemmaculatus*) application to the water toxicity assessment of a peri-urban polluted river of Argentina. *Chemosphere*, 59(4), 577–583. <https://doi.org/10.1016/j.chemosphere.2004.12.039>

De la Torre, F. R., Salibián, A., & Ferrari, L. (1999). Enzyme activities as biomarkers of freshwater pollution: Responses of fish branchial (Na⁺K⁺)-ATPase and liver transaminases. *Environmental Toxicology*, 14(3), 313–319. [https://doi.org/10.1002/\(SICI\)1522-7278\(199907\)14:3<313::AID-TOX4>3.0.CO;2-J](https://doi.org/10.1002/(SICI)1522-7278(199907)14:3<313::AID-TOX4>3.0.CO;2-J)

de la Torre, F. R., Salibián, A., & Ferrari, L. (2007). Assessment of the pollution impact on biomarkers of effect of a freshwater fish. *Chemosphere*, 68(8), 1582–1590. <https://doi.org/10.1016/j.chemosphere.2007.02.033>

- Del Brio, J., Lares, B. A., Parra-Morales, L. B., Sanchez, V. G., Montagna, C. M., & Venturino, A. (2019). Differential detoxifying responses to crude oil water-accommodated fraction in *Hyalalella curvispina* individuals from unpolluted and contaminated sites. *Environmental Toxicology and Pharmacology*, 70, 103191. <https://doi.org/10.1016/j.etap.2019.04.012>
- del Brio, J., Montagna, C. M., Lares, B. A., Parolo, M. E., & Venturino, A. (2018). Chemical characterization and toxicity of water-accommodated fraction of oil on the South American native species *Hyalella curvispina*. *Environmental Toxicology and Pharmacology*, 60, 209–215. <https://doi.org/10.1016/j.etap.2018.04.022>
- Demetrio, P. M., Bonetto, C., & Ronco, A. E. (2014). The Effect of Cypermethrin, Chlorpyrifos, and Glyphosate Active Ingredients and Formulations on *Daphnia magna* (Straus). *Bulletin of Environmental Contamination and Toxicology*, 93(3), 268–273. <https://doi.org/10.1007/s00128-014-1336-0>
- Demetrio, P. M., Bulus Rossini, G. D., Bonetto, C. A., & Ronco, A. E. (2012). Effects of Pesticide Formulations and Active Ingredients on the Coelenterate *Hydra attenuata* (Pallas, 1766). *Bulletin of Environmental Contamination and Toxicology*, 88(1), 15–19. <https://doi.org/10.1007/s00128-011-0463-0>
- Díaz-Jaramillo, M., Laitano, M. V., Gonzalez, M., & Miglioranza, K. S. B. (2018). Spatio-temporal trends and body size differences of OCPs and PCBs in *Laeonereis culveri* (Polychaeta: Nereididae) from Southwest Atlantic estuaries. *Marine Pollution Bulletin*, 136, 107–113. <https://doi.org/10.1016/j.marpolbul.2018.09.008>
- Díaz-Jaramillo, M., Miglioranza, K. S. B., Carriquiriborde, P., Marino, D., Pegoraro, C. N., Valenzuela, G., & Barra, R. (2017). Sublethal effects in *Perinereis gualpensis* (Polychaeta: Nereididae) exposed to mercury-pyrene sediment mixture observed in a multipolluted estuary. *Ecotoxicology*, 26(6), 792–801. <https://doi.org/10.1007/s10646-017-1810-7>
- Díaz-Jaramillo, M., Miglioranza, K. S. B., Gonzalez, M., Barón, E., Monserrat, J. M., Eljarrat, E., & Barceló, D. (2016). Uptake, metabolism and sub-lethal effects of BDE-47 in two estuarine invertebrates with different trophic positions. *Environmental Pollution*, 213, 608–617. <https://doi.org/10.1016/j.envpol.2016.03.009>
- Dorelle, L. S., Da Cuña, R. H., Rey Vázquez, G., Höcht, C., Shimizu, A., Genovese, G., & Lo Nostro, F. L. (2017). The SSRI fluoxetine exhibits mild effects on the reproductive axis in the cichlid fish *Cichlasoma dimerus* (Teleostei, Cichliformes). *Chemosphere*, 171, 370–378. <https://doi.org/10.1016/j.chemosphere.2016.11.141>

- Drewes A.M y Herkovits, J. Eliminación de metales pesados en aguas mediante módulos de fibra hueca: aplicaciones para la industria. *Ingeniería Sanitaria y Ambiental*. 52: 59-62, 2000.
- Eissa, B. L., Ossana, N. A., Ferrari, L., & Salibián, A. (2010). Quantitative Behavioral Parameters as Toxicity Biomarkers: Fish Responses to Waterborne Cadmium. *Archives of Environmental Contamination and Toxicology*, 58(4), 1032–1039. <https://doi.org/10.1007/s00244-009-9434-4>
- Eissa, B. L., Ossana, N. A., Salibián, A., & Ferrari, L. (2017). Assessment of physiological effects of sublethal cadmium on *Cyprinus carpio*. *International Journal of Environment and Health*, 8(2), 139. <https://doi.org/10.1504/IJENVH.2017.083973>
- Eissa, B. L., Salibián, A., & Ferrari, L. (2006). Behavioral Alterations in Juvenile *Cyprinus carpio* (Linnaeus, 1758) Exposed to Sublethal Waterborne Cadmium. *Bulletin of Environmental Contamination and Toxicology*, 77(6), 931–937. <https://doi.org/10.1007/s00128-006-1233-2>
- Embryos: Linkage of Effects to Tissue Residues *Int. J. Environ. Res. Public Health* 4(2), 166-172, 2007
- Fein A, Torchinsky A, Pinchasov M, Katz N, Toder V, Herkovits J. Cadmium embryotoxicity: evidence of a direct effect of cadmium on early rat embryos. *Bull Environ Contam Toxicol*. 59(4):520-4, 1997.
- Ferrari, A., Anguiano, L., Lascano, C., Sotomayor, V., Rosenbaum, E., & Venturino, A. (2008). Changes in the antioxidant metabolism in the embryonic development of the common South American toad *Bufo arenarum* : Differential responses to pesticide in early embryos and autonomous-feeding larvae. *Journal of Biochemical and Molecular Toxicology*, 22(4), 259–267. <https://doi.org/10.1002/jbt.20236>
- Ferrari, A., Anguiano, O. L., Soleño, J., Venturino, A., & Pechen de D'Angelo, A. M. (2004). Different susceptibility of two aquatic vertebrates (*Oncorhynchus mykiss* and *Bufo arenarum*) to azinphos methyl and carbaryl. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 139(4), 239–243. <https://doi.org/10.1016/j.cca.2004.11.006>
- Ferrari, A., Lascano, C. I., Anguiano, O. L., D'Angelo, A. M. P. de, & Venturino, A. (2009). Antioxidant responses to azinphos methyl and carbaryl during the embryonic development of the toad *Rhinella* (*Bufo*) *arenarum* Hensel. *Aquatic Toxicology*, 93(1), 37–44. <https://doi.org/10.1016/j.aquatox.2009.03.003>

- Ferrari, A., Lascano, C., Pechen de D'Angelo, A. M., & Venturino, A. (2011). Effects of azinphos methyl and carbaryl on *Rhinella arenarum* larvae esterases and antioxidant enzymes. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 153(1), 34–39. <https://doi.org/10.1016/j.cbpc.2010.08.003>
- Ferrari, A., Venturino, A., & Pechen de D'Angelo, A. M. (2004). Time course of brain cholinesterase inhibition and recovery following acute and subacute azinphosmethyl, parathion and carbaryl exposure in the goldfish (*Carassius auratus*). *Ecotoxicology and Environmental Safety*, 57(3), 420–425. [https://doi.org/10.1016/S0147-6513\(02\)00069-6](https://doi.org/10.1016/S0147-6513(02)00069-6)
- Ferrari, A., Venturino, A., & Pechén de D'Angelo, A. M. (2007). Effects of carbaryl and azinphos methyl on juvenile rainbow trout (*Oncorhynchus mykiss*) detoxifying enzymes. *Pesticide Biochemistry and Physiology*, 88(2), 134–142. <https://doi.org/10.1016/j.pestbp.2006.10.005>
- Ferrari, A., Venturino, A., & Pechén de D'Angelo, A. M. (2007). Muscular and brain cholinesterase sensitivities to azinphos methyl and carbaryl in the juvenile rainbow trout *Oncorhynchus mykiss*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 146(3), 308–313. <https://doi.org/10.1016/j.cbpc.2007.04.002>
- Ferrari, L., & Salibián, A. (1999). Effect of Cadmium on the Epidermic Structure of *Bufo arenarum* Tadpoles: Influence of the Chemical Composition of the Incubation Media. *Archives of Physiology and Biochemistry*, 107(2), 91–96. <https://doi.org/10.1076/apab.107.2.91.4346>
- Ferrari, L., de la Torre, F. R., Demichelis, S. O., García, M. E., & Salibián, A. (2005). Ecotoxicological assessment for receiving waters with the premetamorphic tadpoles acute assay. *Chemosphere*, 59(4), 567–575. <https://doi.org/10.1016/j.chemosphere.2005.01.045>
- Ferrari, L., Demichelis, S. O., García, M. E., de la Torre, F. R., & Salibián, A. (1997). Premetamorphic anuran tadpoles as test organism for an acute aquatic toxicity assay. *Environmental Toxicology and Water Quality*, 12(2), 117–121. [https://doi.org/10.1002/\(SICI\)1098-2256\(1997\)12:2<117::AID-TOX3>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1098-2256(1997)12:2<117::AID-TOX3>3.0.CO;2-B)
- Ferrari, L., Eissa, B. L., & Salibián, A. (2011). Energy balance of juvenile *Cyprinus carpio* after a short-term exposure to sublethal water-borne cadmium. *Fish Physiology and Biochemistry*, 37(4), 853–862. <https://doi.org/10.1007/s10695-011-9483-2>
- Ferrari, L., Eissa, B. L., Ossana, N. A., & Salibian, A. (2009). Effects of sublethal waterborne cadmium on gills in three teleostean species: scanning electron microscope study.

- Ferrari, L., Gil, D. G., & Vinuesa, J. H. (2011). Breeding and fecundity of the sub-Antarctic crab *Halicarcinus planatus* (Crustacea: Hymenosomatidae) in the Deseado River estuary, Argentina. *Journal of the Marine Biological Association of the United Kingdom*, 91(5), 1023–1029. <https://doi.org/10.1017/S0025315410001840>
- Ferrari, L., Lombardo, R. J., del Giorgio, P., Tortorelli, M. C., & Hernández, D. A. (1989). Effects of formulated ethyl parathion on fertilization of the sea urchin *Pseudechinus magellanicus* (Philippi). *Bulletin of Environmental Contamination and Toxicology*, 42(3), 367–374. <https://doi.org/10.1007/BF01699962>
- Ferreira, M. F., Varela, M. L., Lo Nostro, F., Ansaldo, M., & Genovese, G. (2017). Reproductive aspects of *Notothenia rossii* and *N. coriiceps* (Perciformes, Nototheniidae) at Potter Cove, 25 de Mayo (King George) Island during austral summer. *Polar Biology*, 40(1), 1–11. <https://doi.org/10.1007/s00300-016-1918-x>
- Final Report of the OECD Working Group Meeting on Aquatic Toxicity Testing, pp. 1-77, Copenhagen, Denmark, 29-30 June 1995. Participantes: De Bruijn, J., Kristensen, P., Solbj, J. Bro-Rasmussen F., Graney, R., Hinman, M., Hodson, P., Kallqvist T., Klaschka, U., Morcock, R., Persoone, G., Solomon, K., Tarpkea, M., Tyle, H., Vasseur, P., Herkovits, J. (observer, as a scientists from a no-OECD country), Matthiesen, P., Kopp, H., Mc Cahon, P., Roghair, C., Bias, R., Tiberg, E., Gonzalez-Valero, J., Fleming, R., Hatakeyama, S., Hyne, S., Meyn, O., Stephenson, R., Tsani-Bazaca, E.
- Fogel, M. N., Schneider, M. I., Rimoldi, F., Ladux, L. S., Desneux, N., & Ronco, A. E. (2016). Toxicity assessment of four insecticides with different modes of action on pupae and adults of *Eriopis connexa* (Coleoptera: Coccinellidae), a relevant predator of the Neotropical Region. *Environmental Science and Pollution Research*, 23(15), 14918–14926. <https://doi.org/10.1007/s11356-016-6654-9>
- Fonovich TM, Perez-Coll CS, Fridman O, D'Eramo JL, Herkovits J. Phospholipid changes in *Rhinella arenarum* embryos under different acclimation conditions to copper. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology* 189, 10-16, 2016 <https://doi.org/10.1016/j.cbpc.2016.06.007>
- Fridman O., Aronzon C M, Jorge Herkovits. El impacto del Cadmio sobre la incorporación de Estradiol en embriones de anfibios. En "Salud Ambiental y Humana: Una visión holística" P.85-87 Publicación SETAC. Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

- Fridman O., Corro L. and Herkovits, J. Actividad de 17 β -hidroxiesteroide deshidrogenasa en embriones de Bufo arenarum. Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 124-126, 2003
- Fridman, O., D'Eramo J. L. Herkovits, J., Respuesta Antioxidante en embriones de Bufo arenarum expuestos a radiación ultravioleta (UV-B) en condiciones subletales. Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 144-146, 2003
- Fridman, O.; Corro, L. Herkovits, J. Estradiol uptake, toxicity, metabolism and adverse effect on cadmium treated amphibian embryos. Environmental Health Perspectives 112, 862-866, 2004.
- Frühling, W., Rönnpagel, K., & Ahlf, W. (2001). Effect of zinc and benzalkonium chloride on Nitrosomonas communis and potential nitrification in soil. Environmental Toxicology, 16(5), 439–443. <https://doi.org/10.1002/tox.10002>
- Furland, N. E., Oresti, G. M., Antollini, S. S., Venturino, A., Maldonado, E. N., & Aveldaño, M. I. (2007). Very Long-chain Polyunsaturated Fatty Acids Are the Major Acyl Groups of Sphingomyelins and Ceramides in the Head of Mammalian Spermatozoa. Journal of Biological Chemistry, 282(25), 18151–18161. <https://doi.org/10.1074/jbc.M700709200>
- Gagneten, A. M. (2002). Efectos del herbicida Paraquat sobre el zooplancton. Iheringia. Série Zoologia, 92(3), 47–56. <https://doi.org/10.1590/S0073-47212002000300005>
- Gagneten, A. M., & Paggi, J. C. (2009). Effects of Heavy Metal Contamination (Cr, Cu, Pb, Cd) and Eutrophication on Zooplankton in the Lower Basin of the Salado River (Argentina). Water, Air, and Soil Pollution, 198(1–4), 317–334. <https://doi.org/10.1007/s11270-008-9848-z>
- Gagneten, A. M., Plá, R. R., Regaldo, L., & Paggi, J. C. (2009). Assessment of Bioconcentration Factor of Chromium by Instrumental Neutron Activation Analysis in Argyrodiaptomus falcifer Daday, a Subtropical Freshwater Copepod. Water, Air, and Soil Pollution, 204(1–4), 133–138. <https://doi.org/10.1007/s11270-009-0032-x>
- Gagneten, A. M., Tumini, G., Imhof, A., & Gervasio, S. (2012). Comparative Study of Lead Accumulation in Different Organs of the Freshwater Crab Zilchiopsis oronensis. Water, Air, & Soil Pollution, 223(2), 617–624. <https://doi.org/10.1007/s11270-011-0887-5>

- Galanti, L. N., Amé, M. V., & Wunderlin, D. A. (2013). Accumulation and detoxification dynamic of cyanotoxins in the freshwater shrimp *Palaemonetes argentinus*. *Harmful Algae*, 27, 88–97. <https://doi.org/10.1016/j.hal.2013.05.007>
- Garanzini, D. S., & Menone, M. L. (2015). Azoxystrobin Causes Oxidative Stress and DNA Damage in the Aquatic Macrophyte *Myriophyllum quitense*. *Bulletin of Environmental Contamination and Toxicology*, 94(2), 146–151. <https://doi.org/10.1007/s00128-014-1428-x>
- Garanzini, D. S., Medici, S., Moreyra, L. D., & Menone, M. L. (2019). Acute exposure to a commercial formulation of Azoxystrobin alters antioxidant enzymes and elicit damage in the aquatic macrophyte *Myriophyllum quitense*. *Physiology and Molecular Biology of Plants*, 25(1), 135–143. <https://doi.org/10.1007/s12298-018-0603-7>
- García, M. E., Rodríguez Capítulo, A., & Ferrari, L. (2010). Age-related differential sensitivity to cadmium in *Hyalella curvispina* (Amphipoda) and implications in ecotoxicity studies. *Ecotoxicology and Environmental Safety*, 73(5), 771–778. <https://doi.org/10.1016/j.ecoenv.2009.12.022>
- Genovese, G., Da Cuña, R., Towle, D. W., Maggese, M. C., & Lo Nostro, F. (2011). Early expression of zona pellucida proteins under octylphenol exposure in *Cichlasoma dimerus* (Perciformes, Cichlidae). *Aquatic Toxicology*, 101(1), 175–185. <https://doi.org/10.1016/j.aquatox.2010.09.017>
- Genovese, G., Regueira, M., Da Cuña, R. H., Ferreira, M. F., Varela, M. L., & Lo Nostro, F. L. (2014). Nonmonotonic response of vitellogenin and estrogen receptor α gene expression after octylphenol exposure of *Cichlasoma dimerus* (Perciformes, Cichlidae). *Aquatic Toxicology*, 156, 30–40. <https://doi.org/10.1016/j.aquatox.2014.07.019>
- Genovese, G., Regueira, M., Piazza, Y., Towle, D. W., Maggese, M. C., & Lo Nostro, F. (2012). Time-course recovery of estrogen-responsive genes of a cichlid fish exposed to waterborne octylphenol. *Aquatic Toxicology*, 114–115, 1–13. <https://doi.org/10.1016/j.aquatox.2012.02.005>
- Ghirardi, R., Perotti, M. G., Steciow, M. M., Arellano, M. L., & Natale, G. S. (2011). Potential distribution of *Batrachochytrium dendrobatidis* in Argentina: implications in amphibian conservation. *Hydrobiologia*, 659(1), 111–115. <https://doi.org/10.1007/s10750-010-0270-3>
- Giusto, A., & Ferrari, L. (2008). Copper Toxicity on Juveniles of *Hyalella pseudoazteca* González and Watling, 2003. *Bulletin of Environmental Contamination and Toxicology*, 81(2), 169–173. <https://doi.org/10.1007/s00128-008-9463-0>

- Giusto, A., & Ferrari, L. (2014). Biochemical responses of ecological importance in males of the austral South America amphipod *Hyalella curvispina* Shoemaker, 1942 exposed to waterborne cadmium and copper. *Ecotoxicology and Environmental Safety*, 100, 193–200. <https://doi.org/10.1016/j.ecoenv.2013.11.001>
- Giusto, A., Salibián, A., & Ferrari, L. (2014). Biomonitoring toxicity of natural sediments using juvenile *Hyalella curvispina* (Amphipoda) as test species: evaluation of early effect endpoints. *Ecotoxicology*, 23(2), 293–303. <https://doi.org/10.1007/s10646-013-1173-7>
- Giusto, A., Somma, L. A., & Ferrari, L. (2012). Cadmium toxicity assessment in juveniles of the Austral South America amphipod *Hyalella curvispina*. *Ecotoxicology and Environmental Safety*, 79, 163–169. <https://doi.org/10.1016/j.ecoenv.2011.12.020>
- Gómez, S., Villar, C., & Bonetto, C. (1998). Zinc toxicity in the fish *Cnesterodon decemmaculatus* in the Paraná River and Río de La Plata Estuary. *Environmental Pollution*, 99(2), 159–165. [https://doi.org/10.1016/S0269-7491\(97\)00194-2](https://doi.org/10.1016/S0269-7491(97)00194-2)
- González, C. I., Maine, M. A., Cazenave, J., Hadad, H. R., & Benavides, M. P. (2015). Ni accumulation and its effects on physiological and biochemical parameters of *Eichhornia crassipes*. *Environmental and Experimental Botany*, 117, 20–27. <https://doi.org/10.1016/j.envexpbot.2015.04.006>
- González, C. I., Maine, M. A., Cazenave, J., Sanchez, G. C., & Benavides, M. P. (2015). Physiological and biochemical responses of *Eichhornia crassipes* exposed to Cr (III). *Environmental Science and Pollution Research*, 22(5), 3739–3747. <https://doi.org/10.1007/s11356-014-3558-4>
- Griboff, J., Morales, D., Bertrand, L., Bonansea, R. I., Monferrán, M. V., Asis, R., ... Amé, M. V. (2014). Oxidative stress response induced by atrazine in *Palaemonetes argentinus*: The protective effect of vitamin E. *Ecotoxicology and Environmental Safety*, 108, 1–8. <https://doi.org/10.1016/j.ecoenv.2014.06.025>
- Grondona, S. I., Gonzalez, M., Martínez, D. E., Massone, H. E., & Miglioranza, K. S. B. (2014). Endosulfan leaching from Typic Argiudolls in soybean tillage areas and groundwater pollution implications. *Science of The Total Environment*, 484, 146–153. <https://doi.org/10.1016/j.scitotenv.2014.03.016>
- Guerreño, M., López Armengol, M. F., Luquet, C. M., & Venturino, A. (2016). Comparative study of toxicity and biochemical responses induced by sublethal levels of the pesticide azinphosmethyl in two fish species from North-Patagonia, Argentina. *Aquatic Toxicology*, 177, 365–372. <https://doi.org/10.1016/j.aquatox.2016.06.015>

- Gutierrez, M. F., Gagneten, A. M., & Paggi, J. C. (2010). Copper and Chromium Alter Life Cycle Variables and the Equiproportional Development of the Freshwater Copepod *Notodiaptomus conifer* (SARS). *Water, Air, & Soil Pollution*, 213(1–4), 275–286. <https://doi.org/10.1007/s11270-010-0383-3>
- Gutierrez, M. F., Gagneten, A. M., & Paggi, J. C. (2012). Exposure to sublethal chromium and endosulfan alter the diel vertical migration (DVM) in freshwater zooplankton crustaceans. *Ecotoxicology*, 21(1), 37–47. <https://doi.org/10.1007/s10646-011-0761-7>
- Gutierrez, M. F., Gagneten, A. M., & Paggi, J. C. (2013). Acute and Behavioral Sensitivity of *Mesocyclops longisetus* to Atrazine and Endosulfan Formulations under Predation Pressure. *Water, Air, & Soil Pollution*, 224(1), 1375. <https://doi.org/10.1007/s11270-012-1375-2>
- Gutierrez, M. F., Paggi, J. C., & Gagneten, A. M. (2012). Infodisruptions in predator–prey interactions: Xenobiotics alter microcrustaceans responses to fish infochemicals. *Ecotoxicology and Environmental Safety*, 81, 11–16. <https://doi.org/10.1016/j.ecoenv.2012.04.001>
- Herkovits J, Castañaga LA, D'Eramo JL, Platonova Jourani J. Living organisms influence on environmental conditions: pH modulation by amphibian embryos versus aluminum toxicity. *Chemosphere*, 139, 210-215, 2015
- Herkovits J. Derecho humano al agua y al entorno. Aspectos vinculados a cuencas hídricas. Libro XIV Encuentro Internacional de Derecho Ambiental. Buenos Aires, 2015
- *Herkovits J. Ecotoxicología: Una decision necesaria para el manejo sustentable de cuencas hidricas. *Ingenieria Sanitaria y Ambiental* 125, 25-35, 2015 <http://cuencashidricasecotox.blogspot.com.ar/>
- Herkovits J., Aronzon C. y Pérez Coll C. S. 2009. Exposición de los embriones de anfibio a bajas concentraciones de cobre: respuesta a condiciones de desafíos con el metal, irradiación UV-B y disrupción endócrina. *Química y Toxicología Ambiental en América Latina*, Ed. J.Herkovits, pp 95-97. ISBN 978-987-25370-0-5.
- Herkovits J., Muñoz L.M., Asorey C.M y Lipscomb J. P. Amphitox: un método rápido para detectar la toxicidad remanente al proceso de desinfección del agua de red. En “Salud Ambiental y Humana: Una visión holística” P.34-37 Publicación SETAC. Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1
- Herkovits J., Pérez Coll C. S. y Fridman O. 2009. Acción combinada de estrógenos y cadmio sobre embriones de anfibio: efectos sinérgicos o antagónicos dosis dependientes.

Química y Toxicología Ambiental en América Latina, Ed. J.Herkovits, pp 118-120. ISBN 978-987-25370-0-5.

*Herkovits J.; Rodrigues Capítulo A.; Boyle T.; Servant R.; Pérez-Coll, CS.; Gómez N.; Muñoz L.; Domínguez O.; Cortelezzi A.; Licursi M.; Vanrell T.; Lopez A.; Varela L.; Castañaga, L.; Puszczek E.; Cordero M Estudio ecotoxicológico del arroyo Las Conchitas (Buenos Aires). I Toxicidad en agua y sedimentos. II Parámetros Físico-químicos y relevamiento de la biota.. En "Salud Ambiental y Humana: Una visión holística" pp.50-53. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

Herkovits, J. Bioensayos para Estudios Ecotoxicológicos y de Seguridad Química. Rev. Ing. Sanitaria y Ambiental, 19, 42- 48, 1995.

Herkovits, J. and Helguero, A. L. Copper Toxicity and Copper-Zinc Interactions in Amphibian embryos. The Science of the Total Environment, 221, 1-10, 1998.

Herkovits, J. and Perez-Coll, C. Stage-Dependent Susceptibility of Bufo arenarum embryos to cadmium. Bull. Environ. Contam.Toxicol. 50: 608-611, 1993.

Herkovits, J. and Perez-Coll, C. Amphibian embryos as bioassays to evaluate environmental pollutants and to reduce their toxic effects. Proceedings: Environmental Pollutants, 4th International conference, Barcelona, 623 - 626, 1990.

Herkovits, J. and Perez-Coll, C. Biological effects of Potentized Microdoses. Frontier Perspectives 1, 2, 14 - 15, 1991.

Herkovits, J. and Pérez-Coll, C. Increased resistance against cadmium toxicity by means of pretreatment with low cadmium/zinc concentrations in Bufo arenarum embryos. Biol. Trace Element Res., 49, 171-175, 1995.

Herkovits, J. and Perez-Coll, C. Stage Dependent Uptake of Cadmium in Bufo arenarum Embryos. Bull. Environ. Contam.Toxicol. 56:663-669, 1996.

Herkovits, J. and Perez-Coll, C.S. Antagonism and synergism between lead and zinc in Amphibian larvae.

Herkovits, J. and Perez-Coll, C.S. Zinc protection against delayed development produced by cadmium. Biol. trace Elem. Res. 24: 217-221, 1990.

*Herkovits, J. and Pérez-Coll, CS AMPHITOX: A Standardized Set of Toxicity Tests employing Amphibian Embryos. Its Potential for Customized Hazard Assessment..in Multiple Stressor Effects in Relation to Declining Amphibian Populations. ASTM

(Linder G, Krest S, Sparling D, Little EE, eds) ASTM International. 46-60, 2003. ISBN 0-8031-3464-9

Herkovits, J. Organismos centinela para una gestión ambiental sustentable. Gerencia Ambiental, 2017 Artículo central del Número conmemorativo por los 25 años de la Revista Gerencia Ambiental

Herkovits, J. Paleoecotoxicology: Extending Environmental Toxicology and Chemistry to the Interpretation of the Fossil Record. Environmental Toxicology and Chemistry, 20(8) 1623-1624, 2001.

Herkovits, J. Perez-Coll C. S Ecotoxicología, su importancia para la protección del medio ambiente, la salud humana y la comprensión del proceso evolutivo.. La Revista del ITAES. Vol. 8 N 2 14-29, 2006.

Herkovits, J. Sustentabilidad: Preservación de los servicios del ecosistema. Gerencia Ambiental.. 3, 34, 274-276 y 343, 1997.

Herkovits, J. y Perez-Coll C. Calidad de Agua y vida silvestre: La toxicidad aguda del Níquel y el Aluminio y sus interacciones con el Zinc en embriones de Bufo arenarum. Ingeniería Sanitaria y Ambiental 34, 55-60, 1997.

Herkovits, J. y Perez-Coll, C. Bioensayos para test de toxicidad con embriones de anfibio "ANFITOX". basado en Bufo arenarum. Test Agudo (ANFIAGU), Crónico corto (ANFICOR), Crónico (ANFICRO) y de Estadios Tempranos del Desarrollo (ANFIEMB).. Ingeniería Sanitaria y Ambiental (AIDIS) 42, 24-30 y 43, 50-55, 1999.

Herkovits, J., Herkovits, F.D. and Perez-Coll C.S. Evaluation of Nickel-Zinc Interactions by Means of Bioassays with Amphibian Embryos. Ecotoxicology and Environmental Safety, 45, 266-273, 2000.

Herkovits, J., Herkovits, FD and Perez-Coll, C. Identification of aluminium hazard and aluminium-zinc antagonism in amphibian (Bufo arenarum) embryos.. Environmental Sciences, 5, 1, 57-64, 1997.

Herkovits, J., Paleoecotoxicology: The Impact of Chemical and Physical Stress in the Evolutionary Process. Environmental Health Perspectives, 109 (12) 564-566, 2001.

*Herkovits, J., Perez-Coll, C. and Herkovits, F.D. Ecotoxicological studies of environmental samples from Buenos Aires area using a standardized amphibian embryo toxicity test (AMPHITOX). Environmental Pollution, 116 (1) 177-183, 2002.

*Herkovits, J., Pérez-Coll, C.S., Nievas, G. Y Domínguez, O. Estudio Ecotoxicológico en el Arroyo Claro: Efectos Letales y Subletales. Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 298-301, 2003

Herkovits, J., J. L. D'Eramo and O. Fridman The Effect of UV-B Radiation on Bufo arenarum Embryos Survival and Superoxide Dismutase Activity. International Journal of Environmental Research and Public Health.Vol 3 43-47, 2006.

Herkovits, J., L. Corro, C.S. Perez Coll, O. Domínguez. Fluid motion effect on metal toxicity in Bufo arenarum embryos. Bull. Environ. Contam. Toxicol. 68, 549-554, 2002.

Herkovits, J., Pérez-Coll, CS, Domínguez, O, Servant, R. and C. Vázquez, Exposición a bajas concentraciones de cadmio en embriones de anfibios: resultados de incorporación e incremento en la resistencia a este metal. Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 112-114, 2003

Herkovits, J., Pérez-Coll, CS., Fridman, O., D'Eramo, J.L. y Stockert, J.C. Posibilidades de protección frente al estrés oxidativo producido por agentes físico-químicos en embriones de anfibio. En "Salud Ambiental y Humana: Una visión holística" pp.63-65. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

Herkovits, J.; Cardellini, P., Pavanati, C. y Pérez Coll, C. Cadmium uptake and bioaccumulation in Xenopus laevis embryos at different developmental stages. Ecotoxicology and Environmental Safety, 39, 21-26, 1998.

Herkovits, J.; Cardellini, P.; Pavanati, C. and Perez-Coll, CS. Susceptibility of early life stages of Xenopus laevis to cadmium. Environmental Toxicology and Chemistry, 16, 312-316, 1997.

Herkovits, J.; Muñoz LM.; Asorey, CM. and Lipscomb JC. Application of AMPHITOX assay to determine the toxicity of dichloroacetic and Trichloroacetic acids. Toxicol Sci 72(S-1):156, 2003.

Herkovits, J.; Pérez-Coll, C. and Zeni, S. Protective effect of zinc against spontaneous malformations and lethality in Bufo arenarum embryos. Biol. Trace Elem. Res. 25: 247 - 251, 1989.

*Herkovits, J.; Perez-Coll, CS and Herkovits FD. Ecotoxicity in Reconquista River (Province of Buenos Aires): A preliminary study. Environmental Health Perspectives, 104,2, 186-189, 1996.

- Herkovits, J., Pérez-Coll, C.S., Castañaga, L., Sandoval, M.T., Asorey, C., Muñoz, L., Zuckenberg, C. Los efectos teratogénicos como expresión de calidad ambiental actual y desde una perspectiva evolutiva., Aronzon, C. En "Salud Ambiental y Humana: Una visión holística" pp.88-90. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1
- Hernández, D. A., Lombardo, R. J., Ferrari, L., & Tortorelli, M. C. (1990). Toxicity of ethyl-parathion and carbaryl on early development of sea urchin. *Bulletin of Environmental Contamination and Toxicology*, 45(5), 734–741. <https://doi.org/10.1007/BF01700994>
- Hojman JY., Meichtry J M, Litter M I, Pérez Coll CS. Abatement of toxicity of effluents containing Cr(VI) by heterogeneous photocatalysis. Toxicity assessment by AMPHITOX assay. *Ecotox. Environ. Saf.* 122: 545–550, 2015. DOI:10.1016/j.ecoenv.2015.09.036
- Hued, A. C., Nostro, F. L. Lo, Wunderlin, D. A., & de los Ángeles Bistoni, M. (2013). Reproductive Impairment of a Viviparous Fish Species Inhabiting a Freshwater System with Anthropogenic Impact. *Archives of Environmental Contamination and Toxicology*, 64(2), 281–290. <https://doi.org/10.1007/s00244-012-9826-8>
- Hunt, L., Bonetto, C., Marrochi, N., Scalise, A., Fanelli, S., Liess, M., ... Resh, V. H. (2017). Species at Risk (SPEAR) index indicates effects of insecticides on stream invertebrate communities in soy production regions of the Argentine Pampas. *Science of The Total Environment*, 580, 699–709. <https://doi.org/10.1016/j.scitotenv.2016.12.016>
- Hutler Wolkowicz I., Svartz G., Aronzon C., Pérez Coll C.S." Developmental toxicity of Bisphenol A diglycidyl ether (epoxide resin BADGE) during the early life cycle of a native amphibian species" *Environmental Toxicology and Chemistry* 35 (12): 3031–3038, 2016
- Hutler Wolkowicz, I R., Herkovits J*, and. Pérez Coll C.S (*corresponding author). Stage Dependent Toxicity of Bisphenol A on *Rhinella arenarum* (Anura, Bufonidae) embryos and larvae *Environmental Toxicology* 29, 2, pp. 146 – 154, 2014.
- Hutler Wolkowicz, IR; Aronzon, C., Perez Coll, C.S. Lethal and Sublethal Toxicity of the Industrial Chemical Epichlorohydrin on *Rhinella arenarum* (Anura, Bufonidae) Embryos and Larvae. *J. Hazard. Materials* 263:784-791, 2013
- Iturburu, F. G., Bertrand, L., Mendieta, J. R., Amé, M. V., & Menone, M. L. (2018). An integrated biomarker response study explains more than the sum of the parts: Oxidative stress in the fish *Australoheros facetus* exposed to imidacloprid. *Ecological Indicators*, 93, 351–357. <https://doi.org/10.1016/j.ecolind.2018.05.019>

- Iturburu, F. G., Bertrand, L., Mendieta, J. R., Amé, M. V., & Menone, M. L. (2018). An integrated biomarker response study explains more than the sum of the parts: Oxidative stress in the fish *Australoheros facetus* exposed to imidacloprid. *Ecological Indicators*, 93, 351–357. <https://doi.org/10.1016/j.ecolind.2018.05.019>
- Jergentz, S., Mugni, H., Bonetto, C., & Schulz, R. (2004). Runoff-Related Endosulfan Contamination and Aquatic Macroinvertebrate Response in Rural Basins Near Buenos Aires, Argentina. *Archives of Environmental Contamination and Toxicology*, 46(3). <https://doi.org/10.1007/s00244-003-2169-8>
- Jergentz, S., Mugni, H., Bonetto, C., & Schulz, R. (2005). Assessment of insecticide contamination in runoff and stream water of small agricultural streams in the main soybean area of Argentina. *Chemosphere*, 61(6), 817–826. <https://doi.org/10.1016/j.chemosphere.2005.04.036>
- Jergentz, S., Pessacq, P., Mugni, H., Bonetto, C., & Schulz, R. (2004). Linking in situ bioassays and population dynamics of macroinvertebrates to assess agricultural contamination in streams of the Argentine pampa. *Ecotoxicology and Environmental Safety*, 59(2), 133–141. <https://doi.org/10.1016/j.ecoenv.2004.06.007>
- Lanfranchi, A. L., Menone, M. L., Miglioranza, K. S. B., Janiot, L. J., Aizpún, J. E., & Moreno, V. J. (2006). Striped weakfish (*Cynoscion guatucupa*): A biomonitor of organochlorine pesticides in estuarine and near-coastal zones. *Marine Pollution Bulletin*, 52(1), 74–80. <https://doi.org/10.1016/j.marpolbul.2005.08.008>
- Lanfranchi, A. L., Menone, M. L., Miglioranza, K. S. B., Janiot, L. J., Aizpún, J. E., & Moreno, V. J. (2006). Striped weakfish (*Cynoscion guatucupa*): A biomonitor of organochlorine pesticides in estuarine and near-coastal zones. *Marine Pollution Bulletin*, 52(1), 74–80. <https://doi.org/10.1016/j.marpolbul.2005.08.008>
- Lascano, C. I., Ferrari, A., & Venturino, A. (2011). Sublethal concentrations of azinphos-methyl induce biochemical and morphological alterations in *Rhinella arenarum* embryos. *Chemistry and Ecology*, 27(6), 557–568. <https://doi.org/10.1080/02757540.2011.602971>
- Lascano, C. I., Ferrari, A., Gauna, L. E., Cocca, C., Cochón, A. C., Verrengia, N., & Venturino, A. (2011). Organophosphorus insecticides affect normal polyamine metabolism in amphibian embryogenesis. *Pesticide Biochemistry and Physiology*, 101(3), 240–247. <https://doi.org/10.1016/j.pestbp.2011.10.001>
- Liendro, N., Ferrari, A., Mardirosian, M., Lascano, C. I., & Venturino, A. (2015). Toxicity of the insecticide chlorpyrifos to the South American toad *Rhinella arenarum* at larval

developmental stage. *Environmental Toxicology and Pharmacology*, 39(2), 525–535. <https://doi.org/10.1016/j.etap.2014.12.022>

Lombardo, R. J., Ferrari, L., & Vinuesa, J. H. (1991). Effects of lindane and acetone on the development of larvae of the southern king crab (*Lithodes antarcticus* jaquinot). *Bulletin of Environmental Contamination and Toxicology*, 46(2), 185–192. <https://doi.org/10.1007/BF01691935>

M.E., G., A., R. C., & L., F. (2012). Age differential response of *Hyalella curvispina* to a cadmium pulse: Influence of sediment particle size. *Ecotoxicology and Environmental Safety*, 80, 314–320. <https://doi.org/10.1016/j.ecoenv.2012.03.017>

Mac Loughlin, T. M., Peluso, L., & Marino, D. J. G. (2017). Pesticide impact study in the peri-urban horticultural area of Gran La Plata, Argentina. *Science of The Total Environment*, 598, 572–580. <https://doi.org/10.1016/j.scitotenv.2017.04.116>

Maggioni, T., Hued, A. C., Monferrán, M. V., Bonansea, R. I., Galanti, L. N., & Amé, M. V. (2012). Bioindicators and Biomarkers of Environmental Pollution in the Middle-Lower Basin of the Suquia River (Córdoba, Argentina). *Archives of Environmental Contamination and Toxicology*, 63(3), 337–353. <https://doi.org/10.1007/s00244-012-9785-0>

Marchese, M., Gagneten, A. M., Parma, M. J., & Pavé, P. J. (2008). Accumulation and Elimination of Chromium by Freshwater Species Exposed to Spiked Sediments. *Archives of Environmental Contamination and Toxicology*, 55(4), 603–609. <https://doi.org/10.1007/s00244-008-9139-0>

Mardirosian, M. N., Ceschin, D. G., Lascano, C. I., & Venturino, A. (2017). Molecular effectors in the chronic exposure to arsenic as early and sensitive biomarkers in developing *Rhinella arenarum* toads. *Aquatic Toxicology*, 186, 19–27. <https://doi.org/10.1016/j.aquatox.2017.02.019>

Mardirosian, M. N., Lascano, C. I., Bongiovanni, G. A., & Venturino, A. (2017). Chronic toxicity of arsenic during *Rhinella arenarum* embryonic and larval development: Potential biomarkers of oxidative stress and antioxidant response. *Environmental Toxicology and Chemistry*, 36(6), 1614–1621. <https://doi.org/10.1002/etc.3693>

Mardirosian, M. N., Lascano, C. I., Ferrari, A., Bongiovanni, G. A., & Venturino, A. (2015). Acute toxicity of arsenic and oxidative stress responses in the embryonic development of the common South American toad *Rhinella arenarum*. *Environmental Toxicology and Chemistry*, 34(5), 1009–1014. <https://doi.org/10.1002/etc.2856>

- Mardirosian, M., Bieczynski, F., Luquet, C., Pérez, C. A., Bongiovanni, G., Lascano, C., & Venturino, A. (2017). Arsenic absorption and excretion in chronically exposed developing toad *Rhinella arenarum*. *Environmental Toxicology and Pharmacology*, 52, 255–261. <https://doi.org/10.1016/j.etap.2017.04.014>
- Martínez, D. E., Miglioranza, K. S. B., & Massone, H. E. (2017). Introductory editorial thematic issue: geochemistry of surface processes (III RAGSU). *Environmental Earth Sciences*, 76(2), 96. <https://doi.org/10.1007/s12665-017-6420-4>
- Mastrángelo, M., & Ferrari, L. (2013). *Cnesterodon decemmaculatus* Juveniles as Test Organisms in Toxicity Assessment: Cadmium Case. *Bulletin of Environmental Contamination and Toxicology*, 91(1), 49–54. <https://doi.org/10.1007/s00128-013-0999-2>
- Mastrángelo, M., Dos Santos Afonso, M., & Ferrari, L. (2011). Cadmium toxicity in tadpoles of *Rhinella arenarum* in relation to calcium and humic acids. *Ecotoxicology*, 20(6), 1225–1232. <https://doi.org/10.1007/s10646-011-0667-4>
- Meijide, F. J., Da Cuña, R. H., Prieto, J. P., Dorelle, L. S., Babay, P. A., & Lo Nostro, F. L. (2018). Effects of waterborne exposure to the antidepressant fluoxetine on swimming, shoaling and anxiety behaviours of the mosquitofish *Gambusia holbrooki*. *Ecotoxicology and Environmental Safety*, 163, 646–655. <https://doi.org/10.1016/j.ecoenv.2018.07.085>
- Meijide, F. J., Rey Vázquez, G., Piazza, Y. G., Babay, P. A., Itria, R. F., & Lo Nostro, F. L. (2016). Effects of waterborne exposure to 17 β -estradiol and 4-tert-octylphenol on early life stages of the South American cichlid fish *Cichlasoma dimerus*. *Ecotoxicology and Environmental Safety*, 124, 82–90. <https://doi.org/10.1016/j.ecoenv.2015.10.004>
- Menone, M. L., & Pflugmacher, S. (2005). Effects of 3-chlorobiphenyl on photosynthetic oxygen production, glutathione content and detoxication enzymes in the aquatic macrophyte *Ceratophyllum demersum*. *Chemosphere*, 60(1), 79–84. <https://doi.org/10.1016/j.chemosphere.2004.11.094>
- Menone, M. L., Pesce, S. F., Díaz, M. P., Moreno, V. J., & Wunderlin, D. A. (2008). Endosulfan induces oxidative stress and changes on detoxication enzymes in the aquatic macrophyte *Myriophyllum quitense*. *Phytochemistry*, 69(5), 1150–1157. <https://doi.org/10.1016/j.phytochem.2007.11.016>
- Merlo, C., Abril, A., Amé, M. V., Argüello, G. A., Carreras, H. A., Chiappero, M. S., ... Solís, V. M. (2011). Integral assessment of pollution in the Suquia River (Córdoba, Argentina) as a contribution to lotic ecosystem restoration programs. *Science of The Total Environment*, 409(23), 5034–5045. <https://doi.org/10.1016/j.scitotenv.2011.08.037>

- Merlo, C., Reyna, L., Abril, A., Amé, M. V., & Genti-Raimondi, S. (2014). Environmental factors associated with heterotrophic nitrogen-fixing bacteria in water, sediment, and riparian soil of Suquía River. *Limnologia*, 48, 71–79. <https://doi.org/10.1016/j.limno.2014.06.004>
- Mitton, F. M., Gonzalez, M., Monserrat, J. M., & Miglioranza, K. S. B. (2018). DDTs-induced antioxidant responses in plants and their influence on phytoremediation process. *Ecotoxicology and Environmental Safety*, 147, 151–156. <https://doi.org/10.1016/j.ecoenv.2017.08.037>
- Monferrán, M. V., Galanti, L. N., Bonansea, R. I., Amé, M. V., & Wunderlin, D. A. (2011). Integrated survey of water pollution in the Suquía River basin (Córdoba, Argentina). *J. Environ. Monit.*, 13(2), 398–409. <https://doi.org/10.1039/C0EM00545B>
- Monferran, M. V., Pesce, S. F., Cazenave, J., & Wunderlin, D. A. (2008). Detoxification and antioxidant responses in diverse organs of *Jenynsia multidentata* experimentally exposed to 1,2- and 1,4-dichlorobenzene. *Environmental Toxicology*, 23(2), 184–192. <https://doi.org/10.1002/tox.20326>
- Monserrat, J. M., Garcia, M. L., Ventura-Lima, J., González, M., Ballesteros, M. L., Miglioranza, K. S. B., Wunderlin, D. A. (2014). Antioxidant, phase II and III responses induced by lipoic acid in the fish *Jenynsia multidentata* (Anablápidae) and its influence on endosulfan accumulation and toxicity. *Pesticide Biochemistry and Physiology*, 108, 8–15. <https://doi.org/10.1016/j.pestbp.2013.10.009>
- Monserrat, J. M., Garcia, M. L., Ventura-Lima, J., González, M., Ballesteros, M. L., Miglioranza, K. S. B., Wunderlin, D. A. (2014). Antioxidant, phase II and III responses induced by lipoic acid in the fish *Jenynsia multidentata* (Anablápidae) and its influence on endosulfan accumulation and toxicity. *Pesticide Biochemistry and Physiology*, 108, 8–15. <https://doi.org/10.1016/j.pestbp.2013.10.009>
- Moreyra, L. D., Garanzini, D. S., Medici, S., & Menone, M. L. (2019). Evaluation of Growth, Photosynthetic Pigments and Genotoxicity in the Wetland Macrophyte *Bidens laevis* Exposed to Tebuconazole. *Bulletin of Environmental Contamination and Toxicology*, 102(3), 353–357. <https://doi.org/10.1007/s00128-019-02539-8>
- Mugni, H., Demetrio, P., Bulus, G., Ronco, A., & Bonetto, C. (2011). Effect of Aquatic Vegetation on the Persistence of Cypermethrin Toxicity in Water. *Bulletin of Environmental Contamination and Toxicology*, 86(1), 23–27. <https://doi.org/10.1007/s00128-010-0143-5>
- Mugni, H., Demetrio, P., Marino, D., Ronco, A., & Bonetto, C. (2010). Toxicity Persistence Following an Experimental Cypermethrin and Chlorpyrifos Application in Pampasic

Surface Waters (Buenos Aires, Argentina). Bulletin of Environmental Contamination and Toxicology, 84(5), 524–528. <https://doi.org/10.1007/s00128-010-9986-z>

Mugni, H., Demetrio, P., Paracampo, A., Pardi, M., Bulus, G., & Bonetto, C. (2012). Toxicity Persistence in Runoff Water and Soil in Experimental Soybean Plots Following Chlorpyrifos Application. Bulletin of Environmental Contamination and Toxicology, 89(1), 208–212. <https://doi.org/10.1007/s00128-012-0643-6>

Mugni, H., Paracampo, A., & Bonetto, C. (2013). Nutrient Concentrations in a Pampasic First Order Stream with Different Land Uses in the Surrounding Plots (Buenos Aires, Argentina). Bulletin of Environmental Contamination and Toxicology, 91(4), 391–395. <https://doi.org/10.1007/s00128-013-1079-3>

Mugni, H., Paracampo, A., Demetrio, P., Pardi, M., Bulus, G., Ronco, A., & Bonetto, C. (2016). Toxicity Persistence of Chlorpyrifos in Runoff from Experimental Soybean Plots to the Non-target Amphipod *Hyalella curvispina*: Effect of Crop Management. Archives of Environmental Contamination and Toxicology, 70(2), 257–264. <https://doi.org/10.1007/s00244-015-0187-y>

Mugni, H., Paracampo, A., Demetrio, P., Scalise, A., Solís, M., Fanelli, S., & Bonetto, C. (2015). Acute Toxicity of Endosulfan to the Non-target Organisms *Hyalella curvispina* and *Cnesterodon decemmaculatus*. Bulletin of Environmental Contamination and Toxicology, 95(3), 363–367. <https://doi.org/10.1007/s00128-015-1608-3>

Mugni, H., Paracampo, A., Marrochi, N., & Bonetto, C. (2013). Acute toxicity of cypermethrin to the non target organism *Hyalella curvispina*. Environmental Toxicology and Pharmacology, 35(1), 88–92. <https://doi.org/10.1016/j.etap.2012.11.008>

Mugni, H., Paracampo, A., Solis, M., Fanelli, S., & Bonetto, C. (2014). Acute toxicity of roundup to the nontarget organism *Hyalella curvispina*. Laboratory and field study. Toxicological & Environmental Chemistry, 96(7), 1054–1063. <https://doi.org/10.1080/02772248.2014.993641>

Mugni, H., Ronco, A., & Bonetto, C. (2011). Insecticide toxicity to *Hyalella curvispina* in runoff and stream water within a soybean farm (Buenos Aires, Argentina). Ecotoxicology and Environmental Safety, 74(3), 350–354. <https://doi.org/10.1016/j.ecoenv.2010.07.030>

Natale, G. S., Basso, N. G., & Ronco, A. E. (2000). Effect of Cr(VI) on early life stages of three species of hylid frogs (Amphibia, Anura) from South America. Environmental Toxicology, 15(5), 509–512. [https://doi.org/10.1002/1522-7278\(2000\)15:5<509::AID-TOX21>3.0.CO;2-S](https://doi.org/10.1002/1522-7278(2000)15:5<509::AID-TOX21>3.0.CO;2-S)

- Natale, G. S., Vera-Candioti, J., Ruiz de Arcaute, C., Soloneski, S., Larramendy, M. L., & Ronco, A. E. (2018). Lethal and sublethal effects of the pirimicarb-based formulation Aficida® on *Boana pulchella* (Duméril and Bibron, 1841) tadpoles (Anura, Hylidae). *Ecotoxicology and Environmental Safety*, 147, 471–479. <https://doi.org/10.1016/j.ecoenv.2017.09.007>
- Natale, G., Ammassari, L., Basso, N., & Ronco, A. (2006). Acute and chronic effects of Cr(VI) on *Hypsiboas pulchellus* embryos and tadpoles. *Diseases of Aquatic Organisms*, 72, 261–267. <https://doi.org/10.3354/dao072261>
- Nikoloff, N., Natale, G. S., Marino, D., Soloneski, S., & Larramendy, M. L. (2014). Flurochloridone-based herbicides induced genotoxicity effects on *Rhinella arenarum* tadpoles (Anura: Bufonidae). *Ecotoxicology and Environmental Safety*, 100, 275–281. <https://doi.org/10.1016/j.ecoenv.2013.10.021>
- Nilsen, E., Smalling, K. L., Ahrens, L., Gros, M., Miglioranza, K. S. B., Picó, Y., & Schoenfuss, H. L. (2019). Critical review: Grand challenges in assessing the adverse effects of contaminants of emerging concern on aquatic food webs. *Environmental Toxicology and Chemistry*, 38(1), 46–60. <https://doi.org/10.1002/etc.4290>
- Ocampo, E. H., Menone, M. L., Iturburu, F. G., Nuñez, J. D., & Baeza, J. A. (2014). Effect of the endosymbiotic pea crab *Calyptraeotheres garthi* on the metabolic rate and oxidative status of the slipper limpet *Crepidula cachimilla*. *Invertebrate Biology*, 133(2), 170–179. <https://doi.org/10.1111/ivb.12050>
- Ossana, N. A., Baudou, F. G., Castañé, P. M., Tripoli, L., Soloneski, S., & Ferrari, L. (2019). Histological, Genotoxic, and Biochemical Effects on *Cnesterodon decemmaculatus* (Jenyns 1842) (Cyprinodontiformes, Poeciliidae): Early Response Bioassays to Assess the Impact of Receiving Waters. *Journal of Toxicology*, 2019, 1–13. <https://doi.org/10.1155/2019/4687685>
- Ossana, N. A., Eissa, B. L., Baudou, F. G., Castañé, P. M., Soloneski, S., & Ferrari, L. (2016). Multibiomarker response in ten spotted live-bearer fish *Cnesterodon decemmaculatus* (Jenyns, 1842) exposed to Reconquista river water. *Ecotoxicology and Environmental Safety*, 133, 73–81. <https://doi.org/10.1016/j.ecoenv.2016.06.046>
- Paracampo, A. H., Mugni, H. D., Demetrio, P. M., Pardi, M. H., Bulus, G. D., Asborn, M. D., & Bonetto, C. A. (2012). Toxicity persistence in runoff and soil from experimental soybean plots following insecticide applications. *Journal of Environmental Science and Health, Part B*, 47(8), 761–768. <https://doi.org/10.1080/03601234.2012.676358>
- Paracampo, A., García, I., Mugni, H., Marrochi, N., Carriquiriborde, P., & Bonetto, C. (2015). Fish assemblage of a Pampasic stream (Buenos Aires, Argentina): temporal

variations and relationships with environmental variables. *Studies on Neotropical Fauna and Environment*, 50(3), 145–153.
<https://doi.org/10.1080/01650521.2015.1065658>

Paracampo, A., Solis, M., Bonetto, C., & Mugni, H. (2015). Acute toxicity of chlorpyrifos to the non-target organism *Cnesterodon decemmaculatus*. *International Journal of Environmental Health Research*, 25(1), 96–103.
<https://doi.org/10.1080/09603123.2014.903903>

Parma de Croux, M. J., Loteste, A., & Cazenave, J. (2002). Inhibition of Plasma Cholinesterase and Acute Toxicity of Monocrotophos in a Neotropical Fish *Prochilodus lineatus* (Pisces, Curimatidae). *Bulletin of Environmental Contamination and Toxicology*, 69(3), 356–363. <https://doi.org/10.1007/s00128-002-0070-1>

Peluso J., Aronzon C M., Pérez-Coll C S. Estudio de indicadores fisicoquímicos y toxicológicos del agua superficial del bajo Río Paraná. En “Agua mas Humedales”. p303-318-Abraham E, Quintana R y Mataloni G (ed.) Serie Futuros, UNSAM, Total p. 482

Peluso J., Aronzon C.M., Perez Coll C.S. 2019. "Assessment of environmental quality of waterbodies next to agricultural areas of Buenos Aires Province (Argentina) by means of ecotoxicological studies with *Rhinella arenarum*". *J. Environ. Sci. Health, Part B*. DOI: 10.1080/03601234.2019.1622979 (en prensa, aun inaccesible el DOI)

Peluso, L., Abelando, M., Apartín, C. D., Almada, P., & Ronco, A. E. (2013). Integrated ecotoxicological assessment of bottom sediments from the Paraná basin, Argentina. *Ecotoxicology and Environmental Safety*, 98, 179–186.
<https://doi.org/10.1016/j.ecoenv.2013.09.001>

Peluso, L., Bulus Rossini, G., Salibián, A., & Ronco, A. (2013). Physicochemical and ecotoxicological based assessment of bottom sediments from the Luján River basin, Buenos Aires, Argentina. *Environmental Monitoring and Assessment*, 185(7), 5993–6002. <https://doi.org/10.1007/s10661-012-3000-7>

Pérez Coll C. S., Sandoval M. T. y Herkovits J. 2009. Efectos letales y teratogénicos del Funguicygon® en embriones de *Chaunus* (*Bufo*) *arenarum*. *Química y Toxicología Ambiental en América Latina*, Ed. J.Herkovits, pp 86-88. ISBN 978-987-25370-0-5.

Pérez Coll C. S., Stockert J. C. y Herkovits J. 2009. Análisis de la toxicidad química y fotoquímica de colorantes mediante el bioensayo ANFITOX. *Química y Toxicología Ambiental en América Latina*, Ed. J.Herkovits, pp 92-94. ISBN 978-987-25370-0-5.

Pérez- Coll C.S., Herkovits, J., Domínguez, O. El efecto narcótico producido por sustancias simples y muestras ambientales sobre los embriones de *Bufo arenarum*., Toxicología y Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 173-175, 2003

Pérez Coll CS, Pabón-Reyes C., Meichtry JM, Litter MI. Monitoring of toxicity of As(V) solutions by AMPHITOX test without and with treatment with zerovalent iron nanoparticles. Environ. Toxicol Pharmacol 60: 138–145; 2018. <https://doi.org/10.1016/j.etap.2018.04.018>

Pérez Coll, C.S., Aronzon C.M, and Svartz, G.V. Developmental stages of *Rhinella arenarum* (Anura, Bufonidae) in toxicity studies: AMPHITOX, a customized laboratory assay. Chapter 17, 407-424. In Issues in Toxicology No. 33 “Ecotoxicology and genotoxicology: Non-traditional aquatic models” M. L. Larramendy (Ed) The Royal Society of Chemistry 2017. Published by the Royal Society of Chemistry 2017. Thomas Graham House Science Park, Cambridge, UK

Pérez Coll, C.S., Pabón Reyes, D.C., Meichtry, J.M; Litter, M.I. Lethal and sublethal effects of As(V) on early larval development of *Rhinella arenarum* and utility of AMPHITOX as an ecotoxicological test for As(V) solutions after treatment with zerovalent iron nanoparticles. 363-365 One Century of the Discovery of Arsenicosis in Latin America (1914-2014). As 2014 Arsenic in the Environment Proceedings. Ed: Litter, MI, Nicolli, HB, Meichtry, JM, Quici, N, Bundschuh, J, Bhattacharya, P & Naidu, R CRC Press, 2014

Pérez, Coll, C.S., Fridman, O., D’Eramo, J.L. y Herkovits, J. Las metalotioneínas como biomarcadores de metales en anfibios.. En “Salud Ambiental y Humana: Una visión holística” pp.115-117. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

Pérez, D. J., Lukaszewicz, G., Menone, M. L., & Camadro, E. L. (2011). Sensitivity of *Bidens laevis* L. to mutagenic compounds. Use of chromosomal aberrations as biomarkers of genotoxicity. Environmental Pollution, 159(1), 281–286. <https://doi.org/10.1016/j.envpol.2010.08.028>

Pérez, D. J., Lukaszewicz, G., Menone, M. L., V. Amé, M., & Camadro, E. L. (2014). Genetic and biochemical biomarkers in the macrophyte *Bidens laevis* L. Exposed to a commercial formulation of endosulfan. Environmental Toxicology, 29(9), 1063–1071. <https://doi.org/10.1002/tox.21836>

Pérez, D. J., Menone, M. L., Camadro, E. L., & Moreno, V. J. (2008). Genotoxicity evaluation of the insecticide endosulfan in the wetland macrophyte *Bidens laevis* L.

Pérez, G. L., Torremorell, A., Mugni, H., Rodríguez, P., Vera, M. S., Nascimento, M. do, ... Zagarese, H. (2007). Effects Of The Herbicide Roundup On Freshwater Microbial Communities: A Mesocosm Study. *Ecological Applications*, 17(8), 2310–2322. <https://doi.org/10.1890/07-0499.1>

Pérez, M. R., Rossi, A. S., Bacchetta, C., Elorriaga, Y., Carriquiriborde, P., & Cazenave, J. (2018). In situ evaluation of the toxicological impact of a wastewater effluent on the fish *Prochilodus lineatus*: biochemical and histological assessment. *Ecological Indicators*, 84, 345–353. <https://doi.org/10.1016/j.ecolind.2017.09.004>

Perez-Coll C.S. and Herkovits J. Lethal and teratogenic effects of naringenin evaluated by mean of an amphibian embryo toxicity test (AMPHITOX)" .*Food and Chemical Toxicology* 42, 299-306, 2004.

Pérez-Coll, C. S. , Stockert, JC; Blázquez, A. and Herkovits. The screening of photodynamic toxicity of dyes by means of a bioassay using amphibian embryos. *J Res. J. Chem. Environ.* 11:86-91, 2007.

Pérez-Coll, C.; Herkovits, J. and Salibián, A. Teratogenic effects of cadmium on *Bufo arenarum* during gastrulation. *Experientia*, 42: 1174 - 1176, 1986.

Pérez-Coll, C.; Herkovits, J. and Salibián, A. Embryotoxicity of lead on *Bufo arenarum*. *Bull. Environ. Contam. Toxicol.* 41: 247 - 252, 1988.

Perez-Coll, C.S.; Herkovits, J.; Fridman, O. and D'Eramo, J.L. Metallothioneins and cadmium uptake by the liver in *Bufo arenarum*. *Environ. Pollut.* 97 (3), 311-313, 1997.

Pérez-Coll, CS., Castañaga, L. Domínguez, O. y Herkovits, J. Los efectos de exposiciones a un amplio rango de bajas concentraciones de aluminio sobre los embriones de anfibio. En "Salud Ambiental y Humana: Una visión holística" pp.73-75. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

Pérez-Coll, CS., Domínguez, O., Servant, R. y Herkovits, J. Exposición a bajas concentraciones de níquel: Los efectos o beneficiosos o adversos dependen del pre-tratamiento. . En "Salud Ambiental y Humana: Una visión holística" pp.76-78. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1

Pérez-Coll, C. and Herkovits, J. Stage dependent susceptibility to lead in *Bufo arenarum* embryos. *Environ. Pollut.* 63: 239-245, 1990.

- Pérez-Iglesias, J. M., Franco-Belussi, L., Moreno, L., Tripole, S., de Oliveira, C., & Natale, G. S. (2016). Effects of glyphosate on hepatic tissue evaluating melanomacrophages and erythrocytes responses in neotropical anuran *Leptodactylus latinasus*. *Environmental Science and Pollution Research*, 23(10), 9852–9861. <https://doi.org/10.1007/s11356-016-6153-z>
- Pérez-Iglesias, J. M., Franco-Belussi, L., Natale, G. S., & de Oliveira, C. (2019). Biomarkers at different levels of organisation after atrazine formulation (SIPTRAN 500SC®) exposure in *Rhinella schneideri* (Anura: Bufonidae) Neotropical tadpoles. *Environmental Pollution*, 244, 733–746. <https://doi.org/10.1016/j.envpol.2018.10.073>
- Pérez-Iglesias, J. M., Natale, G. S., Soloneski, S., & Larramendy, M. L. (2018). Are the damaging effects induced by the imazethapyr formulation Pivot® H in *Boana pulchella* (Anura) reversible upon ceasing exposure? *Ecotoxicology and Environmental Safety*, 148, 1–10. <https://doi.org/10.1016/j.ecoenv.2017.10.009>
- Pérez-Iglesias, J. M., Ruiz de Arcaute, C., Natale, G. S., Soloneski, S., & Larramendy, M. L. (2017). Evaluation of imazethapyr-induced DNA oxidative damage by alkaline Endo III- and Fpg-modified single-cell gel electrophoresis assay in *Hypsiboas pulchellus* tadpoles (Anura, Hylidae). *Ecotoxicology and Environmental Safety*, 142, 503–508. <https://doi.org/10.1016/j.ecoenv.2017.04.054>
- Pérez-Iglesias, J. M., Ruiz de Arcaute, C., Nikoloff, N., Dury, L., Soloneski, S., Natale, G. S., & Larramendy, M. L. (2014). The genotoxic effects of the imidacloprid-based insecticide formulation Glacoxan Imida on Montevideo tree frog *Hypsiboas pulchellus* tadpoles (Anura, Hylidae). *Ecotoxicology and Environmental Safety*, 104, 120–126. <https://doi.org/10.1016/j.ecoenv.2014.03.002>
- Pérez-Iglesias, J. M., Soloneski, S., Nikoloff, N., Natale, G. S., & Larramendy, M. L. (2015). Toxic and genotoxic effects of the imazethapyr-based herbicide formulation Pivot H® on monteideo tree frog *Hypsiboas pulchellus* tadpoles (Anura, Hylidae). *Ecotoxicology and Environmental Safety*, 119, 15–24. <https://doi.org/10.1016/j.ecoenv.2015.04.045>
- Pesce, S. F., Cazenave, J., Monferrán, M. V., Frede, S., & Wunderlin, D. A. (2008). Integrated survey on toxic effects of lindane on neotropical fish: *Corydoras paleatus* and *Jenynsia multidentata*. *Environmental Pollution*, 156(3), 775–783. <https://doi.org/10.1016/j.envpol.2008.06.016>
- Pesticide residues in food. Joint FAO/WHO Meeting on Pesticide Residues .Evaluations 1995. PART II Toxicological and environmental. WHO/PCS 96.48. 1-498 (Environmental Monographs, 407-490). Joint meeting of the FAO Panel of Experts

on Pesticide Residues in Food and the Environment and the WHO Core Assessment Group, 1995 (Environmental Core Assessment Group: Calamari, D., Dobson, S., Douben, P.E.T., Herkovits, J., Samsoe-Petersen, L., Temmink, J.H.M.)

- Piazza, Y. G., Pandolfi, M., & Lo Nostro, F. L. (2011). Effect of the Organochlorine Pesticide Endosulfan on GnRH and Gonadotrope Cell Populations in Fish Larvae. *Archives of Environmental Contamination and Toxicology*, 61(2), 300–310. <https://doi.org/10.1007/s00244-010-9621-3>
- Piazza, Y., Pandolfi, M., Da Cuña, R., Genovese, G., & Lo Nostro, F. (2015). Endosulfan affects GnRH cells in sexually differentiated juveniles of the perciform *Cichlasoma dimerus*. *Ecotoxicology and Environmental Safety*, 116, 150–159. <https://doi.org/10.1016/j.ecoenv.2015.03.013>
- Pietsch, C., Wiegand, C., Amé, M. V., Nicklisch, A., Wunderlin, D., & Pflugmacher, S. (2001). The effects of a cyanobacterial crude extract on different aquatic organisms: Evidence for cyanobacterial toxin modulating factors. *Environmental Toxicology*, 16(6), 535–542. <https://doi.org/10.1002/tox.10014>
- R. de la Torre, †, S. O. De, F. (1997). Toxicity of Reconquista River Water: Bioassays with Juvenile *Cnesterodon decemmaculatus*. *Bulletin of Environmental Contamination and Toxicology*, 58(4), 558–565. <https://doi.org/10.1007/s001289900371>
- Rautenberg, G. E., Amé, M. V., Monferrán, M. V., Bonansea, R. I., & Hued, A. C. (2015). A multi-level approach using *Gambusia affinis* as a bioindicator of environmental pollution in the middle-lower basin of Suquía River. *Ecological Indicators*, 48, 706–720. <https://doi.org/10.1016/j.ecolind.2014.09.025>
- Regaldo, L., Gutierrez, M. F., Reno, U., Fernández, V., Gervasio, S., Repetti, M. R., & Gagneten, A. M. (2018). Water and sediment quality assessment in the Colastiné-Corralito stream system (Santa Fe, Argentina): impact of industry and agriculture on aquatic ecosystems. *Environmental Science and Pollution Research*, 25(7), 6951–6968. <https://doi.org/10.1007/s11356-017-0911-4>
- Reno, U., Doyle, S. R., Momo, F. R., Regaldo, L., & Gagneten, A. M. (2018). Effects of glyphosate formulations on the population dynamics of two freshwater cladoceran species. *Ecotoxicology*, 27(7), 784–793. <https://doi.org/10.1007/s10646-017-1891-3>
- Reno, U., Gutierrez, M. F., Regaldo, L., & Gagneten, A. M. (2014). The Impact of Eskoba®, a Glyphosate Formulation, on the Freshwater Plankton Community. *Water Environment Research*, 86(12), 2294–2300. <https://doi.org/10.2175/106143014X13896437493580>

- Rey Vázquez, G., & Lo Nostro, F. (2014). Changes in Hematological Parameters of *Cichlasoma dimerus* (Teleostei, Perciformes) Exposed to Sublethal Concentrations of 4-tert-Octylphenol. *Archives of Environmental Contamination and Toxicology*, 66(3), 463–469. <https://doi.org/10.1007/s00244-014-9997-6>
- Rey Vázquez, G., Meijide, F. J., & Lo Nostro, F. L. (2016). Recovery of the Reproductive Capability Following Exposure to 4-tert-Octylphenol in the Neotropical Cichlid Fish *Cichlasoma dimerus*. *Bulletin of Environmental Contamination and Toxicology*, 96(5), 585–590. <https://doi.org/10.1007/s00128-016-1766-y>
- Rey Vázquez, G., Meijide, F. J., Da Cuña, R. H., Lo Nostro, F. L., Piazza, Y. G., Babay, P. A., Guerrero, G. A. (2009). Exposure to waterborne 4-tert-octylphenol induces vitellogenin synthesis and disrupts testis morphology in the South American freshwater fish *Cichlasoma dimerus* (Teleostei, Perciformes). *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 150(2), 298–306. <https://doi.org/10.1016/j.cbpc.2009.05.012>
- Rimoldi, F., Peluso, L., Bulus Rossini, G., Ronco, A. E., & Demetrio, P. M. (2018). Multidisciplinary approach to a study of water and bottom sediment quality of streams associated with mixed land uses: Case study Del Gato Stream, La Plata (Argentina). *Ecological Indicators*, 89, 188–198. <https://doi.org/10.1016/j.ecolind.2018.01.063>
- Rivera, O.; Belmonte, O. and Herkovits, J. Zinc Protection against cadmium effect on estrual cycle of Wistar rat.. *Biol. trace Elem. Res.* 25: 35 - 38, 1990.
- Rogers E.H., Hunter, E.S. 3rd, Moser VC, Phillips, PM J Herkovits, J Muñoz, L, Hall, LL, Chernoff, N. *J Appl Toxicol.* 25:527-34, 2005.
- Ronco, A., Bulus Rossini, G., Sobrero, C., Apartin, C., Castillo, G., Díaz-Baez, M. C., Mendoza, J. (2005). The application of hazard assessment schemes using the watertox toxicity testing battery. Small-scale freshwater toxicity investigations: Volume 2 - hazard assessment schemes (pp. 233-255) doi:10.1007/1-4020-3553-5_8
- Rosenbaum, E. A., Duboscq, L., Soleño, J., Montagna, C. M., Ferrari, A., & Venturino, A. (2012). Response of biomarkers in amphibian larvae to in situ exposures in a fruit-producing region in North Patagonia, Argentina. *Environmental Toxicology and Chemistry*, 31(10), 2311–2317. <https://doi.org/10.1002/etc.1950>
- Rossi, A., Bacchetta, C., & Cazenave, J. (2017). Effect of thermal stress on metabolic and oxidative stress biomarkers of *Hoplosternum littorale* (Teleostei, Callichthyidae). *Ecological Indicators*, 79, 361–370. <https://doi.org/10.1016/j.ecolind.2017.04.042>

- Ruiz de Arcaute, C., Pérez-Iglesias, J. M., Nikoloff, N., Natale, G. S., Soloneski, S., & Larramendy, M. L. (2014). Genotoxicity evaluation of the insecticide imidacloprid on circulating blood cells of Montevideo tree frog *Hypsiboas pulchellus* tadpoles (Anura, Hylidae) by comet and micronucleus bioassays. *Ecological Indicators*, 45, 632–639. <https://doi.org/10.1016/j.ecolind.2014.05.034>
- Ruiz de Arcaute, C., Salgado Costa, C., Demetrio, P. M., Natale, G. S., & Ronco, A. E. (2012). Influence of existing site contamination on sensitivity of *Rhinella fernandezae* (Anura, Bufonidae) tadpoles to Lorsban®48E formulation of chlorpyrifos. *Ecotoxicology*. <https://doi.org/10.1007/s10646-012-0990-4>
- Salgado Costa, C., Ronco, A. E., Trudeau, V. L., Marino, D., & Natale, G. S. (2018). Tadpoles of the horned frog *Ceratophrys ornata* exhibit high sensitivity to chlorpyrifos for conventional ecotoxicological and novel bioacoustic variables. *Environmental Pollution*, 235, 938–947. <https://doi.org/10.1016/j.envpol.2017.12.096>
- Salibián, A., Minotti, P., Herkovits, J. Effects of low concentrations of ethanol on the embryonic development of *Bufo arenarum*. *Comp. Biochem. Physiol.* 81C, 1: 170 - 174, 1984,
- Sandoval, M.T., Pérez-Coll, C.S. y Herkovits, J. Los efectos del malation sobre los embriones de *Bufo arenarum*: Sobrevida, narcosis, motilidad y teratogénesis. En “Salud Ambiental y Humana: Una visión holística” pp. 82-84. Publicación SETAC, Ed: J. Herkovits, Argentina, 2006. ISBN 13:978-987-05-1059-1
- Sansíñena, J. A., Peluso, L., Salgado Costa, C., Demetrio, P. M., Mac Loughlin, T. M., Marino, D. J. G., Natale, G. S. (2018). Evaluation of the toxicity of the sediments from an agroecosystem to two native species, *Hyalella curvispina* (CRUSTACEA: AMPHIPODA) and *Boana pulchella* (AMPHIBIA: ANURA), as potential environmental indicators. *Ecological Indicators*, 93, 100–110. <https://doi.org/10.1016/j.ecolind.2018.04.061>
- Sfara, V., Fonovich T., Pérez-Coll, C. Sustainable pome fruit production in Patagonia, Argentina. *Technologica Acta* Vol. 9, Nro 2: 57-61, 2016.
- Sobrero, M. C., Rimoldi, F., & Ronco, A. E. (2007). Effects of the Glyphosate Active Ingredient and a Formulation on *Lemna gibba* L. at Different Exposure Levels and Assessment End-Points. *Bulletin of Environmental Contamination and Toxicology*, 79(5), 537–543. <https://doi.org/10.1007/s00128-007-9277-5>
- Soledad Garanzini, D., Gastón Iturburu, F., & Luján Menone, M. (2019). Cambios En La Actividad De Enzimas Antioxidantes De *Myriophyllum Quitense* Expuesta A

Endosulfán A Diferentes Temperaturas. *Revista Internacional de Contaminación Ambiental*, 35(1), 125–135. <https://doi.org/10.20937/RICA.2019.35.01.09>

Solis, M., Arias, M., Fanelli, S., Bonetto, C., & Mugni, H. (2019). Agrochemicals' effects on functional feeding groups of macroinvertebrates in Pampas streams. *Ecological Indicators*, 101, 373–379. <https://doi.org/10.1016/j.ecolind.2019.01.036>

Solis, M., Bonetto, C., Marrochi, N., Paracampo, A., & Mugni, H. (2018). Aquatic macroinvertebrate assemblages are affected by insecticide applications on the Argentine Pampas. *Ecotoxicology and Environmental Safety*, 148, 11–16. <https://doi.org/10.1016/j.ecoenv.2017.10.017>

Solis, M., Mugni, H., Fanelli, S., & Bonetto, C. (2017). Effect of agrochemicals on macroinvertebrate assemblages in Pampasic streams, Buenos Aires, Argentina. *Environmental Earth Sciences*, 76(4), 180. <https://doi.org/10.1007/s12665-017-6476-1>

Solis, M., Mugni, H., Hunt, L., Marrochi, N., Fanelli, S., & Bonetto, C. (2016). Land use effect on invertebrate assemblages in Pampasic streams (Buenos Aires, Argentina). *Environmental Monitoring and Assessment*, 188(9), 539. <https://doi.org/10.1007/s10661-016-5545-3>

Solis, M., Paracampo, A., Bonetto, C., & Mugni, H. (2019). Acute Toxicity of Chlorpyrifos to *Hyalella curvispina*: Comparison of Species Sensitivity and Assessment of Environmental Risk. *Environmental Processes*, 6(1), 107–117. <https://doi.org/10.1007/s40710-019-00352-3>

Sotomayor, V., Chiriotto, T. S., Pechen, A. M., & Venturino, A. (2015). Biochemical biomarkers of sublethal effects in *Rhinella arenarum* late gastrula exposed to the organophosphate chlorpyrifos. *Pesticide Biochemistry and Physiology*, 119, 48–53. <https://doi.org/10.1016/j.pestbp.2015.02.006>

Sotomayor, V., Lascano, C., de D'Angelo, A. M. P., & Venturino, A. (2012). Developmental and polyamine metabolism alterations in *Rhinella arenarum* embryos exposed to the organophosphate chlorpyrifos. *Environmental Toxicology and Chemistry*, 31(9), 2052–2058. <https://doi.org/10.1002/etc.1921>

Stockert JC, Herkovits, J. Photodynamic toxicity and its prevention by antioxidative agents in *Bufo arenarum* embryos. *Toxicology*. 2003 Nov 5;192 (2-3):211-8.

Stockert, J.C. y Herkovits, J., El efecto fotodinámico en embrión de *Bufo arenarum* como modelo de estrés oxidativo: efecto de agentes químicos antioxidantes *Toxicología y*

Química Ambiental. Contribuciones para un Desarrollo Sustentable. SETAC Publication, Ed. Jorge Herkovits, 159-161, 2003

Suárez, R. P., Zaccagnini, M. E., Babbitt, K. J., Calamari, N. C., Natale, G. S., Cerezo, A., Gavier-Pizarro, G. I. (2016). Anuran responses to spatial patterns of agricultural landscapes in Argentina. *Landscape Ecology*, 31(10), 2485–2505. <https://doi.org/10.1007/s10980-016-0426-2>

Svartz G, Aronzon C, Pérez Coll, CS Comparative sensitivity among early life stages of the South American toad to cypermethrin-based pesticide. *Environ. Sci. Pollut. Res.* 23(3):2906-2913, 2016 (DOI) 10.1007/s11356-015-5547-7 ISSN:0944-1344

Svartz G, Marino, D., Ronco, A y Perez Coll, CS Differential uptake of endosulfan in the South American toad under sublethal exposure. *Arch Env.Contam.Toxicol.* 69(1): 104-111, 2015. DOI: 10.1007/s00244-015-0164-5

Svartz G, Papa M, Gosatti M, Jordán M, Soldati A, Samter P, Guraya MM., Pérez Coll C, Perez Catán S. Monitoring the ecotoxicity of γ -Al₂O₃ and Ni/ γ -Al₂O₃ nanomaterials by means of a battery of bioassays. *Ecotoxicol Environ Safety* 144: 200–207, 2017

Svartz G, Sandoval MT; Gosatti M, Perez Catan S. Perez Coll C. 2019 Lethality, neurotoxicity morphological, histological and cellular alterations of Ni-Al nanoceramics on the embryo-larval development of *Rhinella arenarum* *Environ. Toxicol.Pharmacol.* 69:36-43, 2019

Svartz G., Meijide F., Pérez Coll C. Effects of a fungicide formulation on embryo-larval development, metamorphosis, and gonadogenesis of the South American toad *Rhinella arenarum*. *EnvironToxicol Pharmacol* 45: 1–7, 2016

Svartz, G. Pérez Coll, CS Toxicity of endosulfan on embryo-larval development of the South American toad, *Rhinella arenarum* *Environ.Toxicol. Chem.* 33(4):875-881, 2013

Svartz, G. V. and Pérez-Coll, C. S. Comparative toxicity of cypermethrin and a commercial formulation on *Rhinella arenarum* larval development (ANURA: BUFONIDAE) *Int. J. Environ. Health* 6(4):320-329, 2013

Svartz, G. V., Herkovits J*,. Pérez-Coll2 C S (* corresponding author). Sublethal effects of atrazine to embryo-larval development of *Rhinella arenarum* (Anura: Bufonidae). *Ecotoxicology*, 21(4):1251-9. 2012.

Svartz, GV, Aquaroni MM; Pérez Coll CS. Differential sensitivity of developmental stages of the South American toad to a fungicide based on fludioxonil and metalaxyl-M

Environ Sci Pollut Res (2018) 25:23857–23863. <https://doi.org/10.1007/s11356-018-2445-9>

Svartz, GV, Aronzon, CM, Pérez Coll, CS. Combined endosulfan and cypermethrin-induced toxicity to embryo–larval development of *Rhinella arenarum* J. Toxicol. Environ Health Part A: Current issues. 79(5): 197-209, 2016 10.1080/15287394.2015.1126211.

Synergistic Effects of Copper and Butyric Ester of 2,4-Dichlorophenoxyacetic Acid (Esteron Ultra) on Amphibia Embryos. Pérez-Coll, CS. y Herkovits, J. Int. J. Environ. Res. Public Health, 3(4), 343-347, 2006

Sztrum, A A, D'Eramo, JL, and Herkovits, J. Nickel toxicity in embryos and larvae of the South American toad: Effects on cell differentiation, morphogenesis, and oxygen consumption. Environm.Tox. and Chem 30: 5: 1146-1152 2011

Taller Subregional de Sensibilización sobre los Contaminantes Orgánicos Persistentes (COPs) en el Período Perinatal: Compuestos tipo Dioxinas. Herkovits, J. Reunión UNEP, realizado en Puerto Iguazú (Argentina) 1-3 de Abril, 1998

The Toxic Effects of Xenobiotics on the Health of Humans and Animals. Editors: Yanzhu Zhu, Alex Boye, Mathilde Body-Malapel, and Jorge Herkovits, 2017 Hindawi.

Tolussi, C. E., Gomes, A. D. O., Kumar, A., Ribeiro, C. S., Lo Nostro, F. L., Bain, P. A., Moreira, R. G. (2018). Environmental pollution affects molecular and biochemical responses during gonadal maturation of *Astyanax fasciatus* (Teleostei: Characiformes: Characidae). Ecotoxicology and Environmental Safety, 147, 926–934. <https://doi.org/10.1016/j.ecoenv.2017.09.056>

Troncoso, I. C., Cazenave, J., Bacchetta, C., & Bistoni, M. de los Á. (2012). Histopathological changes in the gills and liver of *Prochilodus lineatus* from the Salado River basin (Santa Fe, Argentina). Fish Physiology and Biochemistry, 38(3), 693–702. <https://doi.org/10.1007/s10695-011-9551-7>

uyón, N. F., Roggio, M. A., Amé, M. V., Wunderlin, D. A., & Bistoni, M. A. (2016). Dynamics of Aromatase and Physiological Indexes in Male Fish as Potential Biomarkers of Anthropogenic Pollution. Bulletin of Environmental Contamination and Toxicology, 97(5), 632–638. <https://doi.org/10.1007/s00128-016-1925-1>

Valdés, M. E., Amé, M. V., Bistoni, M. de los A., & Wunderlin, D. A. (2014). Occurrence and bioaccumulation of pharmaceuticals in a fish species inhabiting the Suquía River basin (Córdoba, Argentina). Science of The Total Environment, 472, 389–396. <https://doi.org/10.1016/j.scitotenv.2013.10.124>

- Varela, M. L., Ferreira, M. F., Da Cuña, R. H., Lo Nostro, F. L., Genovese, G., & Meijide, F. J. (2017). Dynamics of ovarian maturation throughout the reproductive cycle of the Neotropical cichlid fish *Cichlasoma dimerus* (Teleostei, Cichliformes). *Canadian Journal of Zoology*, 95(7), 485–498. <https://doi.org/10.1139/cjz-2016-0198>
- Venturino, A. (2017). Toxicogenomics: new strategies for ecotoxicology studies in autochthonous species I. A glade in the entangled path towards the “fingerprint” of environmental impact *International Journal of Environment and Health*, 8(3), 203. <https://doi.org/10.1504/IJENVH.2017.086179>
- Venturino, A., Anguiano, O. L., Gauna, L., Cocca, C., Bergoc, R. M., & Pechen de D’Angelo, A. M. (2001). Thiols and polyamines in the potentiation of malathion toxicity in larval stages of the toad *Bufo arenarum*. *Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology*, 130(2), 191–198. [https://doi.org/10.1016/S1532-0456\(01\)00241-1](https://doi.org/10.1016/S1532-0456(01)00241-1)
- Venturino, A., Gauna, L. E., Bergoc, R. M., & Pechen de D’Angelo, A. M. (1992). Effect of exogenously applied polyamines on malathion toxicity in the toad *Bufo arenarum* Hensel. *Archives of Environmental Contamination and Toxicology*, 22(1), 135–139. <https://doi.org/10.1007/BF00213312>
- Venturino, A., Gauna, L., Bergoc, R. M., & Pechen de D’Angelo, A. M. (2001). Toxicokinetics of Malathion in Larval Stages of the Toad *Bufo arenarum* (Hensel): Effect of Exogenous Spermidine. *Pesticide Biochemistry and Physiology*, 70(3), 142–150. <https://doi.org/10.1006/pest.2001.2550>
- Venturino, A., Montagna, C. M., & Pechen de D’Angelo, A. M. (2007). Risk assessment of magnacide® herbicide at río colorado irrigation channels (argentina). Tier 3: studies on native species. *Environmental Toxicology and Chemistry*, 26(1), 177. <https://doi.org/10.1897/06-085R.1>
- Venturino, A., Rosenbaum, E., Caballero De Castro, A., Anguiano, O. L., Gauna, L., Fonovich De Schroeder, T., & Pechen De D’Angelo, A. M. (2003). Biomarkers of effect in toads and frogs. *Biomarkers*, 8(3–4), 167–186. <https://doi.org/10.1080/1354700031000120116>
- Vera Candiotti, J., Natale, G. S., Soloneski, S., Ronco, A. E., & Larramendy, M. L. (2010). Sublethal and lethal effects on *Rhinella Arenarum* (Anura, Bufonidae) tadpoles exerted by the pirimicarb-containing technical formulation insecticide Aficida®. *Chemosphere*, 78(3), 249–255. <https://doi.org/10.1016/j.chemosphere.2009.10.064>
- Vera, M. S., Lagomarsino, L., Sylvester, M., Pérez, G. L., Rodríguez, P., Mugni, H., Pizarro, H. (2010). New evidences of Roundup® (glyphosate formulation) impact on the



periphyton community and the water quality of freshwater ecosystems. *Ecotoxicology*, 19(4), 710–721. <https://doi.org/10.1007/s10646-009-0446-7>

Vilches, C., Giorgi, A., Mastrángelo, M., & Ferrari, L. (2011). Non-point Contamination Homogenizes the Water Quality of Pampean Streams. *Bulletin of Environmental Contamination and Toxicology*, 87(2), 147–151. <https://doi.org/10.1007/s00128-011-0312-1>

Vreys, N., Amé, M. V., Filippi, I., Cazenave, J., Valdés, M. E., & Bistoni, M. A. (2019). Effect of Landscape Changes on Water Quality and Health Status of *Heptapterus mustelinus* (Siluriformes, Heptapteridae). *Archives of Environmental Contamination and Toxicology*, 76(3), 453–468. <https://doi.org/10.1007/s00244-018-00593-7>

Yanzhu Zhu, Alex Boye, Mathilde Body-Malapel, Jorge Herkovits The Toxic Effects of Xenobiotics on the Health of Humans and Animals, Editorial (2 pages), Article ID 4627872, Volume 2017

Zambrano, M. J., Rautenberg, G. E., Bonifacio, A. F., Filippi, I., Amé, M. V., Bonansea, R. I., & Hued, A. C. (2018). Effects of water quality on aspects of reproductive biology of *Cnesterodon decemmaculatus*. *Science of The Total Environment*, 645, 10–21. <https://doi.org/10.1016/j.scitotenv.2018.07.084>